

THE FIRST SURVEY OF THE NORTHERN SKY FOR TeV POINT SOURCES

K. Wang,^{1,a} R. Atkins,² W. Benbow,³ D. Berley,⁴ M.-L. Chen,^{4,b} D. G. Coyne,³ B. L. Dinges,² D. E. Dorfan,³ R. W. Ellsworth,⁵ A. Falcone,⁶ L. Fleysher,⁷ R. Fleysher,⁷ G. Gisler,⁸ J. A. Goodman,⁴ T. J. Haines,⁸ C. M. Hoffman*,⁸ S. Hugenberger,⁹ L. A. Kelley,³ I. Leonor,⁹ M. McConnell,⁶ J. F. McCullough,³ J. E. McEnery,² R. S. Miller,⁶ A. I. Mincer,⁷ M. F. Morales,³ P. Nemethy,⁷ J. M. Ryan,⁶ B. Shen,¹ A. Shoup,⁹ G. Sinnis,⁸ A. J. Smith,⁴ G. W. Sullivan,⁴ O. T. Tumer,¹ M. O. Wascko,¹ S. Westerhoff,^{3,c} D. A. Williams,³ T. Yang,³ G. B. Yodh⁹ (The Milagro Collaboration)

¹University of California, Riverside, CA 92521

²University of Wisconsin, Madison, WI 53706

³University of California, Santa Cruz, CA 95064

⁴University of Maryland, College Park, MD 20742

⁵George Mason University, Fairfax, VA 22030

⁶University of New Hampshire, Durham, NH 03824-3525

⁷New York University, New York, NY 10003

⁸Los Alamos National Laboratory, Los Alamos, NM 87545

⁹University of California, Irvine, CA 92717

^aNow at Fresher Information Corp., San Francisco, CA 94105

^bNow at Brookhaven National Laboratory, Upton, NY 11973

^cNow at Columbia University, New York, NY 10027

Abstract

The development of imaging atmospheric Cherenkov telescopes has led to the detection of several astronomical point sources of TeV photons. These sources had first all been observed previously at other wavelengths. Only a very small fraction of the sky has been studied in the TeV region. We report here the results of the first search for steady TeV point sources anywhere in the northern sky, made with the Milagro air-shower-particle detector from February 1997 to May 1998.

PACS:

Keywords: Gamma-ray astronomy

* Corresponding author: Cyrus Hoffman MS H803 Los Alamos National Laboratory,
Los Alamos, NM 87544 USA Tel.: +1 505 667 5876;
e-mail: cy@lanl.gov.

1. Introduction

Many objects in the universe emit electromagnetic radiation via thermal processes. High-energy gamma rays can only be produced in non-thermal, energetic acceleration processes or more exotic phenomena such as evaporating primordial black holes, topological defects, and neutralino annihilation. Observations of high-energy gamma rays have begun to reveal a picture of high-energy acceleration processes in the universe. Several reviews of high-energy gamma-ray astronomy have recently appeared.[1,2,3]

Cosmic gamma rays up to several GeV can be directly detected with satellite-based detectors. The EGRET instrument [4] surveyed the sky above 100 MeV and detected diffuse Galactic and extragalactic emission, five pulsars, at least 66 blazars, and about 170 "unidentified" objects found along the Galactic plane.[5] Many of the blazars exhibit time variability, while the pulsars and many of the other sources seem to be steady. The unidentified sources appear to be of two classes: relatively faint mid-latitude sources that may be associated with the Gould belt, and brighter low-latitude sources that may be Galactic pulsars or supernova remnants.[6] A catalog of sources above 1 GeV contains 57 sources, including 10 that were not identified in the lower-energy catalogs.[7] This catalog includes the five pulsars and 21 blazars seen at lower energies, as well as 30 unidentified sources near the Galactic plane. The sensitivity of EGRET above 1 GeV is too low to permit any study of transient sources.

Gamma-ray fluxes fall rapidly with increasing energy so that even the brightest source is too faint to be observed above ~ 10 GeV in the relatively small detectors that can be placed in satellites: thus ground-based detectors, which can have much larger collection areas, are used at higher energies. A high-energy gamma ray interacts high in the earth's atmosphere producing an extensive air shower (EAS). Ground-based gamma-ray telescopes detect the EAS products that survive to ground level.

Imaging atmospheric Cherenkov telescopes (ACT) have been successfully employed over the last decade in the energy region from ~ 400 GeV to 10 TeV. ACTs detect Cherenkov light produced in the atmosphere by the ultra-relativistic charged particles in the EAS. ACTs have detected emission from a number of sources including three pulsar-driven nebulae (the Crab, Vela, and PSR1706-44), and three blazars (Markarian 421, Markarian 501, and 1ES2344+514). The TeV emission from the pulsar-driven nebulae appears to be steady, while the blazars exhibit strong flares. However, because of their small field of view (~ 10 msr) and low duty factor (typically 10%), ACTs are not well suited to scan the sky for TeV sources. In fact, only a small fraction of the sky has been examined with ACTs.

Extensive air-shower particle detector arrays (EAS arrays) have been widely used to search for gamma-ray sources above ~ 50 TeV. An EAS array has typically consisted of a sparse array of scintillation counters that detects a small fraction of the charged EAS particles that reach ground level. An EAS array can operate 24 hours per day regardless of weather, and can simultaneously observe the entire overhead sky; an EAS array is able to observe every source in its field of view every day of the year. No convincing evidence for steady gamma-ray emission above 50 TeV from anywhere in the Northern sky has been obtained with these detectors. [8]

We report here on the results of the first systematic search of the northern sky for steady point sources of TeV emission using data from the Milagrito detector, an EAS array sensitive to gamma rays in the 1-TeV region.

2. The Milagrito Detector

The Milagrito detector [9], which operated from February 1997 to May 1998, was the first EAS array to use a large volume of water as the detection medium. Milagrito used photomultiplier tubes (PMTs) deployed under 1-2 m of water to detect the Cherenkov radiation produced in the water by relativistic charged EAS particles. Most of the shower particles (including the gamma rays, which produce electrons and positrons via pair production) falling within the area of the water can be detected with this technique. Thus Milagrito had good sensitivity to gamma rays at and below 1 TeV, while maintaining the large-aperture and high-duty-factor attributes of an EAS array.

Milagrito utilized 228 8"-diameter PMTs located in a plane on a 2.8 m x 2.8 m grid near the bottom of a large, covered pond of water. Data were taken with 1, 1.5, and 2 m of water above the PMTs. The pond is located at an altitude of 2650 m above sea level (atmospheric overburden = 750 g/cm²) in the Jemez mountains west of Los Alamos, New Mexico. As in a conventional EAS array, the direction of the primary gamma ray was reconstructed by measuring the relative times at which the individual PMTs were struck by light produced by particles in the shower front. To reduce the impact of stray light, only tubes registering pulses larger than 2 photoelectrons are used in the shower fits.

The angular resolution for cosmic-ray events was studied by comparing the incident directions separately reconstructed with two independent interleaved portions of the detector. The resulting angular resolution is in good agreement with expectations from a detailed Monte Carlo simulation.[9] However, this study is not sensitive to certain systematic effects such as those due to errors in the reconstructed shower core location. Observations of the cosmic ray shadow of the moon [10] indicate that the simulation reproduces the actual detector point spread function quite well. These observations are for events initiated by cosmic rays, and it is the point spread function for events initiated by gamma rays that must be known to optimize the search for a point source of gamma rays. The simulations indicate that the angular resolution is 10-20% better for gamma-ray events. This is in general agreement with the shape of the signal from Markarian 501 observed with Milagrito.[11] The technique used to search for point-source emission, described below, depends only weakly on the detailed shape of the detector point spread function. Further details on Milagrito are available in Ref. [9].

3. All-Sky Survey Technique

Milagrito had no ability to distinguish showers initiated by gamma rays from background showers initiated by charged cosmic rays. Because of the large number of isotropic background cosmic-ray showers, a strong point gamma-ray source would appear as an excess number of showers coming from a small region of the sky, consistent with the Milagrito point spread function. The analysis in this study required $N_{\text{Fit}} > 50$, where N_{Fit} is the number of PMTs participating in the fit to the shower plane. A total of 3.4×10^9 events are used in this analysis. "Rectangular" sky bins rather than circular bins are used because they can be easily configured to cover the entire sky. With the $N_{\text{Fit}} > 50$ cut, the optimal bin on the sky in which to search for a

point source is $\sim 2^\circ \times 2^\circ$. The actual bin size in celestial coordinates is 2.22° [declination, δ] \times $2.22^\circ/\cos(\delta)$ [right ascension, RA] with one abnormally-sized bin (at each δ) ending at RA = 360° . Because the expected significance for a source depends only weakly on small changes of the bin size, [11] a bin a little larger than optimal was chosen to reduce the trials factor in the survey. A map consisting of a grid of 4319 bins covers the sky from $\delta = -3.9^\circ$ to $\delta = 71.7^\circ$. Because an unknown source may be located near an edge of a bin, three additional maps were also examined: one with bins shifted by 1.11° in δ , one with bins shifted by $1.11^\circ/\cos(\delta)$ in RA, and one with bins shifted by these amounts in both δ and RA. Thus the survey covers the region from $\delta = -5^\circ$ to $\delta = 71.7^\circ$.

An accurate estimate of the expected background is needed in each bin. The expected background in a given bin depends upon its exposure and on the detector efficiency expressed in local coordinates (δ , and hour angle, HA). The background estimation in each bin is determined using the data itself.[12] Random "background" events are generated for each observed event by associating the event local coordinates (δ , HA) with times selected randomly from all event times recorded over a 2-hr period. New values of celestial coordinates are then calculated for each background event. Because the selection of a new event time amounts to a rotation of the celestial sphere with respect to the earth, this method changes only the RA of an event but not its δ . Fifteen background events are generated for each actual event so that the statistical error for the estimated background is small. The background events are then binned in the same manner as the data events: the expected background level is the number of background events falling into a source bin, divided by 15. It has been shown that this method produces background events that track changes in the trigger rate and are free of systematic errors.[12] Once signal and background maps have been produced, they are compared on a bin-by-bin basis and the statistical significance of the observed number of events being due to a fluctuation of the expected background is calculated using the method of Li and Ma.[13]

4. Results

Figure 1a shows the distribution of the statistical significance of the observed number of events above the expected cosmic-ray background for each of the 4319 independent bins of the first map. None of these bins shows a significant excess (or deficit), considering the number of independent bins examined. The distributions for the shifted maps are shown in Figures 1b-d. The bins in a given map are independent of each other but they are not independent of the bins in the other maps. A Monte Carlo study indicates that the effective trials factor for a large excess to appear anywhere in a study of all of the maps is $\approx 2.5 \times 4319 \approx 10,800$. The largest observed excess is 3.9σ , corresponding to a post-trials probability of $\sim 50\%$ of being due to a fluctuation of the background. Thus no strong steady source of TeV emission is observed in the northern sky for the period from February 1997 to May 1998.

The sensitivity of Milagrito has been evaluated using the Monte Carlo simulation.[9] This has been verified by comparing the observed rate of gamma rays from Markarian 501 from February-October 1997[11] with the flux measured by several air Cherenkov telescopes,[14] and by comparing the observed rate of cosmic-ray events with the measured flux.[15] The upper limit on the number of gamma-ray events in a bin varies according to the number of observed events in that bin. Rather than give gamma-ray flux upper limits on a bin-by-bin basis, we give upper

limits for a typical bin as a function of declination. The upper limit on the flux from a source at a given declination, Φ_γ , expressed in terms of the upper limit on the number of gamma-ray events, N_γ , for the source bin at that declination is:

$$N_\gamma = \int \Phi_\gamma(E) A_{\text{EFF}}(E, \delta, \text{HA}(t)) \epsilon \, dE \, dt ,$$

where $A_{\text{EFF}}(E, \delta, \text{HA})$ is the effective area of Milagrito as a function of gamma-ray energy and the local coordinates of the source position, $\text{HA}(t)$ is the hour angle of the source as a function of time, and ϵ is the fraction of events from a source that fall within the source bin. The time integral is taken over the lifetime of the observations. Assuming that a differential source spectrum is $\propto E^{-2.5}$, the resulting typical 90% confidence-level upper limit on the integral gamma-ray flux above 1 TeV for any steady source as a function of declination is shown in Figure 2. For comparison, the integral gamma-ray flux above 1 TeV from the Crab, measured by the Whipple collaboration [16], is also shown in Figure 2.

We have also searched for steady emission from the four known northern TeV sources, the Crab, Mrk 421, Mrk 501, and 1ES2344+514. In each case, the bin in any of the maps that is most nearly centered on the actual source position was used. The results are shown in Table 1. Because no bin is exactly centered on the source position and the bin used in this analysis is larger than the optimal bin, a small loss of sensitivity is to be expected. The observed excess for the Crab (1.1σ) is in good agreement with the expected signal size (1.3σ), considering the expected 7% loss of sensitivity. The excess from Mrk 501 (2.8σ) is less than reported in Ref. 11 (3.7σ). This is partially due to the small expected loss in sensitivity discussed above (an expected loss of $\sim 12\%$), and partially due to statistical fluctuations in the number of events passing the tighter N_{Fit} cut and in the signal size in the larger bin used here.

Source	RA	Dec.	$\Delta(\text{RA})$	$\Delta(\text{Dec.})$	#on	#excess	Significance
Crab	83.6°	22.0°	-0.02°	0.33°	9.15 x 10 ⁵	1050	1.1 σ
Mrk 421	166.1°	38.2°	0.51°	-0.13°	1.13 x 10 ⁶	-850	-0.8 σ
Mrk 501	253.5°	39.8°	0.40°	0.31°	1.15 x 10 ⁶	2980	2.8 σ
1ES2344+514	356.2°	51.4°	0.04°	-0.24°	1.06.x 10 ⁶	227	0.2 σ

Table I: Observations with Milagrito of the four known northern TeV sources. $\Delta(\text{RA})$ [$\Delta(\text{Dec.})$] is the angular distance from the center of the best bin to the actual source location in right ascension [declination]. #n is the number of observed events in the source bin, and #excess is the difference between #on and the estimated number of background events.

5. Conclusions and Outlook

The first systematic survey of the northern sky ($-5^\circ < \delta < 72^\circ$) for point TeV sources has been made with data from the Milagrero air shower detector. No source with a steady flux larger than $\sim 5x$ the gamma-ray flux from the Crab has been observed for the period from February 1997 - May 1998. A paper describing a search of the northern sky for episodic emission on a variety of time scales is in preparation.[17]

Milagrito was dismantled in 1998 to install the Milagro detector. Milagro has better sensitivity than Milagrero due to its larger size, the use of reflecting conical baffles on each photomultiplier tube to increase its light-collection area and eliminate late light due to horizontally-traveling light, and the ability to distinguish photon-initiated showers from background showers with a second layer of PMTs. Milagro began data taking in December 1999. An array of 180 4.5-m² water detectors is being installed over an area of 10,000 m² surrounding the pond, which should further improve the sensitivity of Milagro. One of the goals of Milagro is to sensitively survey the northern sky for TeV gamma-ray sources over a wide variety of time scales.

ACKNOWLEDGMENTS

We acknowledge the contributions of the many people who helped construct Milagrero. We especially thank R. S. Delay and M. Schneider, who were indispensable in the construction, maintenance and operation of Milagrero. This work was supported in part by the National Science Foundation, The U. S. Department of Energy Office of High Energy Physics, The U. S. Department of Energy Office of Nuclear Physics, Los Alamos National Laboratory, the University of California, the Institute of Geophysics and Planetary Physics, The Research Corporation, and the California Space Institute.

REFERENCES

- [1] Ong, R., *Physics Reports*, **305**, 93 (1998).
- [2] Hoffman, C. M., C. Sinnis, P. Fleury, and M. Punch, *Rev. Mod. Physics* **71**, 897 (1999).
- [3] Weekes, T. C. *Physica Scripta*, **T85**, 195 (2000).
- [4] Thompson, D. J., *et al.* *Ap. J. Suppl. Ser.* **86**, 629 (1993).
- [5] Thompson, D. J., *et al.* *Ap. J. Suppl. Ser.* **101**, 259 (1995); Hartman, R. C., *et al.* *Ap. J. Suppl. Ser.* **123**, 79 (1999).
- [6] Gehrels, N., D. J. Macomb, D. L. Bertsch, D. J. Thompson and R. C. Hartman, *Nature*, **404**, 363 (2000).
- [7] Lamb, R. C., and D. J. Macomb, *Ap. J.* **488**, 872 (1997).
- [8] Alexandreas, D. E., *et al.* *Ap. J.* **383**, L53 (1991); McKay, T. A., *et al.*, *Ap. J.* **417**, 742 (1993).
- [9] Atkins, R., *et al.*, *Nucl. Instrum. Meth. Phys. Res.* **449A**, 478 (2000).
- [10] Wascko, M., *et al.*, *Proceedings of the XXVI International Cosmic Ray Conference, Salt Lake City*, **7**, 218 (1999); and Wascko, M., *et al.*, in preparation.
- [11] Atkins, R., *et al.*, *Ap. J.* **525**, L25 (1999).
- [12] The technique of estimating the background is described more fully in Alexandreas, D. E., *et al* Ref. [8], and Alexandreas, D. E. *et al.*, *Nucl. Instrum. Meth. Phys. Res. A* **328**, 570 (1993).
- [13] Li, T. and Y. Ma, *Ap. J.* **272**, 317 (1983).
- [14] Samuelson, F. W. *et al*, *ApJ*, **501**, L17 (1998); Aharonian, F., *et al*, *A&A*, **342**, 69 (1999).

- [15] Asakimori, K. *et al.*, Ap. J. **502** 278 (1998); and Wiebel-Sooth, B., P. L. Biermann, and H. Meyer, Astron. & Astrop. **330**, 389 (1998).
- [16] Hillas, A. M., *et al.*, Ap J. **503**, 744 (1998).
- [17] McCullough, J. *et al.*, in preparation.




Fig. 1a. - The statistical significances for the observed number of events above the expected cosmic-ray background for each of the 4319 independent bins of the first sky map. The smooth curve is a fit to a Gaussian.

Fig. 1b-d. - The statistical significances for the bins of the three shifted maps.




Fig. 2.- The typical 90% confidence level upper limit on the integral flux above 1 TeV for a steady source as a function of declination (δ). The triangle shows the integral flux above 1 TeV from the Crab (from Ref. 14) for comparison.