

Appendix A: Answers to Selected Problems

Ch 1: Units and Problem Solving

- | | | | |
|-----|--|-----|---|
| 1a. | A person of height 5 ft. 11 in. is 1.80 m tall | 12. | 196 cm ² |
| 1b. | The same person is 180 cm | 13. | 250 cm ³ |
| 2a. | 3 seconds = 1/1200 hours | 14. | 8:1, each side goes up by 2 cm, so it will change by 2 ³ |
| 2b. | 3×10^3 ms | 15. | 3.5×10^{51} :1 |
| 3. | 87.5 mi/hr | 16. | 72,000 km/h |
| 4c. | if the person weighs 150 lb. this is equivalent to 668 N | 17. | 0.75 kg/s |
| 5. | Pascals (Pa), which equals N/m ² | 18. | 8×2^N cm ³ /sec; N is for each second starting with 0 seconds for 8 cm ³ |
| 6. | 168 lb., 76.2 kg | 19. | About 12 million |
| 7. | 5 mi/hr/s | 20. | About 1½ trillion (1.5×10^{12}) |
| 8. | 15.13 m | 21. | [a] = N/kg = m/s ² |
| 9. | 11.85 m | | |
| 10. | 89,300 mm | | |
| 11. | 2025 mm ² | | |

Ch 2: Energy Conservation

- | | | | |
|-----|---|------|--|
| 1. | d | 8a. | 34 m/s at B; 28 m/s at D, 40 m/s at E, 49 m/s at C and F; 0 m/s at H |
| 2. | (discuss in class) | 8b. | 96 m |
| 3a. | 5.0×10^5 J | 9a. | 1.7 J |
| 3b. | 3.7×10^5 J | 9b. | 1.3 m/s |
| 3c. | Chemical bonds in the food. | 9c. | 0.4 J, 0.63 m/s |
| 3d. | 70 m/s | 10a. | 1.2 m/s ² |
| 4a. | 5.0×10^5 J | 10b. | 130 J |
| 4b. | 108 m/s | 11a. | 6750 J |
| 5a. | 450,000 J | 11b. | 2.25×10^5 J |
| 5b. | 22,500 J | 11c. | 1.5×10^5 J/gallon of gas |
| 5c. | 16,875 J | 12. | 0.76 m |
| 5d. | 21.2 m/s | | |
| 5e. | 9.18 m | | |
| 7b. | KE = 504,600 J; U _g = 1,058,400 J;
E _{total} = 1,563,000 J | | |

Ch 3: One-Dimensional Motion (answers assume 9.8 m/s² for acceleration of gravity)

- | | | | |
|-----|---|------|---|
| 2. | b | 8. | 6 minutes |
| 3. | a,c,d | 9d. | 20 meters |
| 4. | a,d | 9e. | 40 meters |
| 5a. | Zyan | 9f. | 2.67 m/s |
| 5b. | Ashaan is accelerating because the distance he travels every 0.1 seconds is increasing, so the speed must be increasing | 9g. | 6 m/s |
| 5c. | Ashaan | 9h. | Between t = 15 s and t = 20 sec because your position goes from x = 30 m to x = 20 m. |
| 5d. | Zyan | 9i. | You went forwards and backwards |
| 5f. | Ashaan | 10a. | 7.7 m/s ² |
| | | 10b. | 47 m, 150 feet |

- 10c. 34 mph
 11a. 1.22 m
 11b. 4.9 m/s
 11c. 2.46 m/s
 11d. -4.9 m/s
 12b. 1 second
 12c. at 2 seconds
 12d. 4m
 13a. 250 m
 13b. 16 m/s, -16m/s
 13c. 14s for round trip
 14. 6 times higher
 15. -31m/s^2
 16a. 23 m/s
- 16b. 3.6 seconds
 16c. 28 m
 16d. 45m
 17a. 25 m/s
 17b. 30 m
 17c. -10 m/s^2
 18. 2 m/s^2
 19a. $v_0 = 0$
 19b. The acceleration of gravity (9.8 m/s^2)
 19c. The acceleration of gravity (9.8 m/s^2)
 19d. 60 m
 20a. 0.3 m/s^2
 20b. 0.5 m/s

Ch 4: Two-Dimensional and Projectile Motion (answers assume 9.8 m/s^2 for acceleration of gravity)

- 7a. 13 m
 7b. 41 degrees
 7c. $v_y = 26\text{ m/s}; v_x = 45\text{ m/s}$
 7d. 56 degrees, 14 m/s
 9. 32 m
 10a. 0.5 s
 10b. 0.8 m/s
 11. 105 m
 12. $t = 0.60\text{ s}$, 1.8 m below target
 13. 28 m.
 14a. 3.5 s.
 14b. 35 m; 15 m
15. 40 m; 9.1 m
 16. 1.3 seconds, 7.1 meters
 17. 50 m; $v_{0y} = 30\text{ m/s}$; 50° ; on the way up
 18. 4.4 s
 19. 19°
 20. 0.6 s
 21. 2.3 m/s
 22. 6 m
 23. 1.4 seconds
 24a. yes
 24b. 18 m/s @ -46 degrees from horizontal
 25. 22 m/s @ 62 degrees

Ch 5: Newton's Laws

4. Zero; weight of the hammer minus the air resistance.
 5. 2 forces
 6. 1 force
 7. No
 8. The towel's inertia resists the acceleration
 9a. Same distance
 9b. You go farther
 9c. Same amount of force
 11a. 98 N
 11b. 98 N
 13. 32 N
 14. 5.7 m/s^2
 17. $F_x = 14\text{ N}, F_y = 20\text{ N}$
 18. Left picture: $F = 23\text{N}$ 98° , right picture:
 $F = 54\text{ N}$ 5°
 19. 3 m/s^2 east
 20. 4 m/s^2 ; 22.5° NE
 21. 0.51
 22. 0.2
 23. The rope will not break because his weight of 784 N is distributed between the two ropes.
24. Yes, because his weight of 784 N is greater than what the rope can hold.
 25. Mass is 51 kg and weight is 82 N
 26a. While accelerating down
 26b. 686 N
 26c. 826 N
 27a. 390 N
 27b. 490 N
 28. 0.33
 29. 3.6 kg
 30. $g\sin\theta$
 31b. 20 N
 31c. 4.9 N
 31d. 1.63 kg
 31e. Eraser would slip down the wall
 32a. 1450 N
 32b. 5600 N
 32c. 5700 N
 32d. Friction between the tires and the ground
 32e. Fuel, engine, or equal and opposite reaction
 33b. 210 N
 33c. Normal Force decreases as angle increases.

- 33d. 2.8 m/s^2
 33e. 28 m/s
 33f. no
 33g. 69 N
 33h. 57 N
 33i. 40 N
 33j. 0.33
 33k. 0.09
 35a. zero
 35b. $-kx_0$
 36b. $f_1 = \mu_k m_1 g \cos\theta; f_2 = \mu_k m_2 g \cos\theta$
 36c. Ma
- 36d. $T_A = (m_1 + m_2)(a + \mu \cos\theta)$
 and $T_B = m_2 a + \mu m_2 \cos\theta$
 36e. Solve by using $d = 1/2at^2$ and substituting h for d
 37a. Yes, because it is static and you know the angle and m_1
 37b. Yes, T_A and the angle gives you m_1 and the angle and T_C gives you m_2 , $m_1 = T_A \cos 25^\circ/g$ and $m_2 = T_C \cos 30^\circ/g$
 38a. 3 seconds
 38d. 90 m
 42a. $1.5 \text{ N}; 2.1 \text{ N}; 0.71$

Ch 6: Centripetal Forces

- 5a. 100 N
 5b. 10 m/s^2
 6a. 25 N towards her
 6b. 25 N towards you
 7a. 14.2 m/s^2
 7b. $7.1 \times 10^3 \text{ N}$
 7c. friction between the tires and the road
 8. $.0034g$
 9a. $6.2 \times 10^5 \text{ m/s}^2$
 9b. The same as a.
 10. $3.56 \times 10^{22} \text{ N}$
 11. $4.2 \times 10^{-7} \text{ N}$; very small force
 12. $g = 9.8 \text{ m/s}^2$; you'll get close to this number but not exactly due to some other small effects
 13a. $4 \times 10^{26} \text{ N}$
 13b. gravity
 13c. $2 \times 10^{41} \text{ kg}$
 14. $.006 \text{ m/s}^2$
 15a. .765
- 15b. 4880 N
 16a. $\sim 10^{-8} \text{ N}$ very small force
 16b. Your pencil does not accelerate toward you because the frictional force on your pencil is much greater than this force.
 17a. $4.23 \times 10^7 \text{ m}$
 17b. $6.6 R_e$
 17d. The same, the radius is independent of mass
 18. $1.9 \times 10^7 \text{ m}$
 19. You get two answers for r , one is outside of the two stars one is between them, that's the one you want, $1.32 \times 10^{10} \text{ m}$ from the larger star.
 22a. $v = 28 \text{ m/s}$
 22b. v-down, a-right
 22c. f-right
 22d. Yes, 640 N

Ch 7: Momentum Conservation

8. 37.5 m/s
 9. $v_1 = 2v_2$
 10a. $24 \frac{\text{kg}\cdot\text{m}}{\text{s}}$
 10b. 0.364 m/s
 10c. $22 \frac{\text{kg}\cdot\text{m}}{\text{s}}$
 10d. 109 N
 10e. 109 N due to Newton's third law
 11. $2.0 \text{ kg}, 125 \text{ m/s}$
 12. 21 m/s to the left
 13. 3250 N
 14a. 90 sec
 14b. $1.7 \times 10^5 \text{ sec}$
 15a. 60 m/s
 15b. $.700 \text{ sec}$
 15c. yes, 8.16 m
16. 0.13 m/s to the left
 17a. 11000 N to the left
 17b. tree experienced same average force of 11000 N but to the right
 17c. 2500 lb.
 17d. about $2.5 \text{ "g"} \text{ s}$ of acceleration
 18a. no change
 18b. the last two cars
 19a. 0.0057 s
 19b. 2.85 kg
 20a. 0.0058 m/s^2
 20b. 3.5 m/s^2
 21a. 15 m/s
 21b. 49° S of E
 22b. 1.72 m/s 8.36°

Ch 8: Energy & Force

- | | | | |
|------|---|------|---|
| 6a. | $7.18 \times 10^9 \text{ J}$ | 19. | 8 m/s same direction as the cue ball and 0 m/s |
| 6b. | 204 m/s | 20. | $v_{\text{golf}} = -24.5 \text{ m/s}; v_{\text{pool}} = 17.6 \text{ m/s}$ |
| 7a. | 34 m/s @ B; 28 m/s @ D; 40 m/s @ E; 49 m/s @ C and F; 0 m/s @ H | 21. | 2.8 m |
| 7b. | 30 m | 22a. | 0.57 m/s |
| 7c. | Yes, it makes the loop!! | 22b. | Leonora's |
| 8a. | 2.3 m/s | 22c. | 617 J |
| 8c. | No, the baby will not clear the hill. | 23a. | 19.8 m/s |
| 9a. | 29,500 J | 23b. | 8.8 m/s |
| 9b. | 13 m | 23c. | 39.5 m |
| 11a. | 86 m | 24a. | 89 kW |
| 11b. | 220 m | 24b. | 0.4 |
| 12a. | 48.5 m/s | 24c. | 15.1 m/s |
| 12b. | 128 N | 25. | 43.8 m/s |
| 13. | 0.32 m/s each | 29a. | $3.15 \times 10^5 \text{ J}$ |
| 14a. | 10 m/s | 29b. | 18.0 m/s |
| 14b. | 52 m | 29c. | 2.41 m |
| 15a. | $1.1 \times 10^4 \text{ N/m}$ | 29d. | 7900 J |
| 15b. | 2 m above the spring | 30a. | $v_0/14$ |
| 16. | 96% | 30b. | $mv_0^2/8$ |
| 18a. | .008 m | 30c. | $7mv_0^2/392$ |
| 18b. | 5.12° | 30d. | 71% |

Ch 9: Rotational Motion

- | | | | |
|------|---|------|---|
| 2a. | $9.74 \times 10^{37} \text{ kg m}^2$ | 10b. | 40 N |
| 2b. | $1.33 \times 10^{47} \text{ kg m}^2$ | 10c. | 0.02 rad/s ² |
| 2c. | 0.5 kg m^2 | 10d. | 25 s |
| 2d. | 0.28 kg m^2 | 11a. | Coin with the hole |
| 2e. | 0.07 kg m^2 | 11b. | Coin with the hole |
| 3. | a. True, all rotate 2π for 86,400 sec which is 24 hours, | 12a. | weight |
| 3b. | True, $\omega = 2\pi/t$ and $t=86,400 \text{ s}$ | 12b. | 19.6 N |
| 3f. | True, L is the same | 12c. | plank's length (0.8m) left of the pivot |
| 3g. | $L = I\omega$ and $I = 2/5 mr^2$ | 12d. | 15.7 N m, |
| 3h. | True, $K = \frac{1}{2} I\omega^2$ & $I = 2/5 mr^2$ sub-in $K = 1/5 mr^2\omega^2$ | 12e. | Ba. weight, Bb. 14.7 N, Bc. plank's length (0.3m) left of the pivot, Bd. 4.4 N m, Ca. weight, Cb. 13.6 N, Cc. plank's length (1.00 m) right of the pivot, Cd. 13.6 N m, f) 6.5 N m CC, g) no, net torque doesn't equal zero |
| 3i. | True, $K = \frac{1}{2} I\omega^2$ & $I = mr^2$ sub-in $K = 1/2 mr^2\omega^2$, | 13a. | $7.27 \times 10^{-6} \text{ Hz}$ |
| 4a. | 250 rad | 13b. | 7.27 Hz |
| 4b. | 40 rad | 14a. | 100 Hz |
| 4c. | 25 rad/s | 14b. | $1.25 \times 10^5 \text{ J}$ |
| 4d. | Force applied perpendicular to radius allows α | 14c. | 2500 J-s |
| 4e. | 0.27 kg m^2 , | 14d. | 12,500 m-N |
| 4f. | $K_5 = 84 \text{ J}$ and $K_{10} = 340 \text{ J}$ | 15. | $1.92 \text{ rads/sec} = 0.306 \text{ rev/s}$ |
| 6. | Moment of inertia at the end $1/3 ML^2$ at the center $1/12 ML^2$, angular momentum, $L = I\omega$ and torque, $\tau = I\alpha$ change the in the same way | 16. | 2300 N |
| .8. | Lower | 17b. | 771 N, 1030 N |
| 9. | Iron ball | 17c. | 554 kgm ² |
| 10a. | 200 N team | 17d. | 4.81 rad/sec^2 |
| | | 18a. | 300 N |
| | | 18b. | 240N, -22 N |

- 18c. .092
 19a. -1.28 mN
 19b. CCW
 20a. 2280 N
 20b. 856 N toward beam, 106 N down
 20c. 425 kg m^2
 20d. 3.39 rad/sec^2
 21a. 1411 kg
 21c. 17410 N
 22a. -2.4 m/s^2
 22b. 1.25 seconds

Ch 10: Simple Harmonic Motion

- 1a. Buoyant force and gravity
 1b. $T = 6 \text{ s}$, $f = 1/6 \text{ Hz}$
 2a. $9.8 \times 10^5 \text{ N/m}$
 2b. 0.5 mm
 2c. 22 Hz, no,
 3. $3.2 \times 10^3 \text{ N/m}$
 4a. 110 N/m
 4d. $v(t) = (25)\cos(83t)$
 7a. 0.0038 s
 7b. 0.0038 s
 10. 4 times
 11. 0.04 m
 12a. 16 Hz
 12b. 16 complete cycles but 32 times up and down, 315 complete cycles but 630 times up and down
 12c. 0.063 s
 13a. 24.8 J, 165 N, 413 m/s^2
 13b. 11.1 m/s, 0, 0
 13c. 6.2 J, 18.6 J, 9.49 m/s, 82.5 N, 206 m/s^2
 13d. .169 sec, 5.9 Hz
 14b. .245 J
 14c. 1.40 m/s
 14d. 1.00 m/s
 14f. 2.82 N
 14g. 3.10 N

Ch 11: Wave Motion and Sound

1. 390 Hz
 2a. 4 Hz
 2b. It was being driven near its resonant frequency.
 2c. 8 Hz, 12 Hz
 2d. (Note that earthquakes rarely shake at more than 6 Hz).
 5a. 7 nodes including the 2 at the ends
 5b. 3.6 Hz
 6. 1.7 km
 7a. 1.7 cm
 7b. 17 m
 8a. $4.3 \times 10^{14} \text{ Hz}$
 8b. $2.3 \times 10^{-15} \text{ s}$ - man that electron is moving fast
 9a. 2.828 m
 9b. 3.352 m
 9c. $L = \frac{1}{4} \lambda$ so it would be difficult to receive the longer wavelengths.
 10. Very low frequency
 11b. Same as closed at both ends
 13. 1.9 Hz or 2.1 Hz.
 14. 0.53 m
 15. 2.2 m, 36 Hz; 1.1 m, 73 Hz; 0.733 m, 110 Hz; 0.55 m, 146 Hz
 16. 430 Hz ; $1.3 \times 10^3 \text{ Hz}$; $2.1 \times 10^3 \text{ Hz}$; $3.0 \times 10^3 \text{ Hz}$;
 17a. The tube closed at one end will have a longer fundamental wavelength and a lower frequency.
 17b. If the temperature increases the wavelength will not change, but the frequency will increase accordingly.
 18. struck by bullet first.
 19. 80 Hz; 0.6 m
 20a. 0.457 m
 20b. 0.914 m
 20c. 1.37 m
 21. 2230 Hz; 2780 Hz; 2970 Hz
 22. 498 Hz
 23. 150 m/s

Ch 12: Electricity

- 11b. 1350 N
11c. 1350 N
12a. $1.1 \times 10^9 \text{ N/C}$
12b. 9000 N
13. $F_g = 1.0 \times 10^{-47} \text{ N}$ and $F_e = 2.3 \times 10^{-8} \text{ N}$. The electric force is 39 orders of magnitudes bigger.
14. $1.0 \times 10^{-4} \text{ C}$
16a. down
16b. Up 16c. $5.5 \times 10^{11} \text{ m/s}^2$
16e. $2.9 \times 10^8 \text{ m/s}^2$
17a. Toward the object
17b. $3.6 \times 10^4 \text{ N/C}$ to the left with a force of $2.8 \times 10^{-7} \text{ N}$
18. Twice as close to the smaller charge, so 2 m from $12\mu\text{C}$ charge and 1 m from $3\mu\text{C}$ charge.
19. 0.293 N and at 42.5°
20. 624 N/C and at an angle of -22.4° from the + x-axis.
21a. 7500V
21b. 1.5 m/s
22a. $6.4 \times 10^{-17} \text{ N}$
22b. 1300V
22c. $2.1 \times 10^{-16} \text{ J}$
22d. $2.2 \times 10^7 \text{ m/s}$
23b. 0.25m
23c. $F_T = 0.022 \text{ N}$
23d. $0.37 \mu\text{C}$

Ch 13: Electric Circuits – Batteries and Resistors

- 1a. 4.5C
1b. 2.8×10^{19} electrons
2a. 0.11 A
2b. 1.0 W
2c. 2.5×10^{21} electrons
2d. 3636 W
3a. 192Ω
3b. 0.42 W
4a. 5.4 mV
4b. $1.4 \times 10^{-8} \text{ A}$
4c. $7.3 \times 10^{-11} \text{ W}$, not a lot
4d. $2.6 \times 10^{-7} \text{ J}$
5. left = brighter, right = longer
6a. 224 V
6b. 400 W by 100Ω and 48 W by 12Ω
6c. 448 W
7b. 8.3Ω
8. 0.5A
10. 0.8A and the 50Ω on the left
11a. 0.94 A
11b. 112 W
11c. 0.35 A
11d. 0.94 A
11e. 50, 45, 75 Ω
- 11f. dimmer; total resistance increases
11g. 45Ω
12a. 0.76 A
12b. 7.0 W
13b. 1000 W
14a. lightbulb A
14b. lightbulbs B,C
14c. i. All; ii. C; iii. None
14d. i. A dimmer, D brighter, C out
ii. A dimmer, B,C brighter
15a. 9.1Ω
15b. 29.1Ω
15c. 10.8Ω
15d. 26.8Ω
15e. 1.8A
15f. 21.5V
15g. 19.4V
15h. 6.1V
15i. 0.24A
15j. 16 W hr
16a. 3.66Ω
16b. 0.36A
16c. 1.32 V
23a. 10V

Ch 14: Magnetism

1. No: if $v = 0$ then $F = 0$; yes: $F = qE$
4a. Into the page
4b. Down the page
4c. Right
5. Both pointing away from north
8. 7.6 T, south
9. Down the page; 60 N
10a. To the right, $1.88 \times 10^4 \text{ N}$
- 10b. 91.7 m/s
10c. It should be doubled
11. East $1.5 \times 10^4 \text{ A}$
12. 0.00016 T; if CCW motion, B is pointed into the ground.
13. $1.2 \times 10^5 \text{ V}$, counterclockwise
14a. 15 V
14b. Counter-clockwise

- 15a. 2×10^{-5} T
 15b. Into the page
 15c. 2.8 N/m
 15d. CW
 16a. 2.42×10^8 m/s
 16b. 9.69×10^{-12} N
 16c. .0055 m
 17. E/B
 18a. 8×10^{-7} T
 18b. 1.3×10^{-6} C
 19a. 0.8 V
 19b. CCW
 19c. .064 N
 19d. .16 N/C
 19e. .13 w
 20a. 1.11×10^8 m/s
 20b. 9.1×10^{-30} N << 6.4×10^{-14} N
- 20d. .00364 T
 20e. .173 m
 20f. 7.03×10^{16} m/s²
 20g. 3.27°
 21. 19.2 V
 22a. 8.39×10^7 m/s
 22b. 2.68×10^{-13} N, -y
 22c. 2.95×10^{17} m/s²
 22d. .00838 m
 22e. 1.68×10^6 N/C
 22f. 16,800 V
 23a. 1.2×10^{-6} T, +z
 23b. 1.5×10^{-17} N, -y
 23c. 96 N/C, -y

Ch 15: Electric Circuits—Capacitors

- 2a. 4×10^7 V
 2b. 4×10^9 J
 4a. 100 V
 4b. A greater voltage created a stronger electronic field, or as charges build up they repel each other from the plate.
 5. 21 V, note that V is squared here
 6a. 3.3 F
 6b. 54Ω
 7a. 200 V
 7b. 5×10^{-9} F
 7c. 2.5×10^{-9} F
 8a. 6V
 8b. 0.3A
 8c. 18V
 8d. 3.6×10^{-4} C
 8e. 3.2×10^{-3} J
 8f. i) 80μF ii) 40μF iii) 120μF
 9a. 26.7μF
- 9b. $166.7 \mu\text{F}$
 10a. 9.0×10^{-3} N/C
 10b. 1.4×10^{-15} N
 10c. 1.6×10^{15} m/s²
 10d. 3.3×10^{-11} s
 10e. 8.9×10^{-7} m
 10f. 5.1×10^{-30}
 11. discuss in class
 12. discuss in class
 13. 10 V
 14a. 0.43 seconds
 14b. 1.8 seconds
 14c. 4.7×10^{-4} C
 14d. No, it will asymptotically approach it
 14e. It will rise rapidly and tail off approaching asymptote of 12 V
 14f. $V_{\text{res}} = 12\text{V} - \text{voltage across capacitor}$
 15. $128 \text{ k}\Omega$

Ch 16: Electric Circuits—Advanced

- 1a. 4.9×10^{-5} H
 1b. -9.8×10^{-5} V
 2. Zero
 3a. Yes
 3b. No
 3c. Because they turn current flow on and off.
 4a. 0.5 V
 4b. 0.05 A
 4c. 0.05 A
 4d. 5.5 V
 4e. 8.25V
- 4f. $11 \times$
 5a. On
 5b. On
 5c. On, on, off, on, off, off, on, on
 6b. $10.9 \mu\text{F}$
 6c. 195Ω
 6d. 169Ω
 6e. 1.39 A
 6f. -42°
 6g. 115Hz

Ch 17: Light

- | | | | |
|------|--|------|----------------------------|
| 3. | 2200 blue wavelengths | 19e. | -1 |
| 4. | 65000 x-rays | 20a. | 6 units |
| 5. | 6×10^{14} Hz | 20b. | bigger; $M = 3$ |
| 6. | 3.3 m | 21c. | 1.5 units |
| 8b. | vacuum & air | 21d. | $\frac{2}{3}$ |
| 8c. | 1.96×10^8 m/s | 22c. | 21/4 units |
| 9. | 6.99×10^{-7} m; 5.26×10^{-7} m | 22e. | $-\frac{2}{3}$ |
| 12. | Absorbs red and green. | 23c. | 5.3 units |
| 13. | 25° | 25b. | 22.5 mm |
| 15. | 33.3° | 27. | 32 cm |
| 16a. | 49.7° | 28a. | 10.2° |
| 16b. | No such angle | 28b. | 27 cm |
| 16c. | 48.8° | 28c. | 20 cm |
| 17b. | 11.4 m | 29a. | 0.72 m |
| 17c. | 11.5 m | 31. | 54 cm, 44cm, 21 cm, 8.8 cm |
| 18. | 85 cm | 33. | 13.5° |
| 19c. | 4 units | 34. | 549 nm |

Ch 18: Fluids

- | | | | |
|------|---|------|--|
| 1. | 0.84 | 12. | .0081 |
| 2. | 1.4×10^5 kg | 13a. | 12500 J/m^3 |
| 3a. | 90% of the berg is underwater | 13b. | 184 kPa |
| 3b. | 57% | 13c. | 1.56 kW |
| 4b. | 5.06×10^4 N | 13d. | 2.57 kW |
| 4c. | 7.05 m/s^2 | 13e. | 11.8 A |
| 5. | 4.14 m/s | 13f. | \$12.6 |
| 6. | 40 coins | 14a. | 611 kPa |
| 7b. | upward | 14b. | 6 atm |
| 7c. | 4.5 m/s^2 | 15b. | 500,000 N |
| 7d. | Cooler air outside, so more initial buoyant force | 16a. | 27 m/s^2 , (2.7 g) upward |
| 7e. | Thin air at high altitudes weighs almost nothing, so little weight displaced. | 16b. | 1600 N |
| 8a. | At a depth of 10 cm, the buoyant force is 2.9 N | 16c. | 2200 N |
| 8d. | The bottom of the cup is 3 cm in radius | 17a. | 10 N |
| 9a. | 83,000 Pa | 17b. | 10.5 N |
| 9b. | 104 N | 17c. | 11 N |
| 9c. | 110 N | 17d. | 11 N |
| 10a. | 248 kPa | 18a. | “The Thunder Road” |
| 10b. | 591 kPa | 18b. | 2.0 m (note: here and below, you may choose differently) |
| 10c. | 1081 kPa | 18c. | 33.5 m^3 |
| | | 18e. | 3.5 million N |
| | | 18f. | 111 MPa |

Ch 19: Thermodynamics and Heat Engines

- | | | | |
|------|-----------------------------|------|---|
| 18. | 517 m/s | 24a. | No |
| 19. | 1.15×10^{12} K | 24b. | allowed by highly improbable state. More likely states are more disordered. |
| 21. | 40 N | 25a. | 8.34×10^{23} |
| 22. | $\approx 10^{28}$ molecules | 25b. | 6.64×10^{-27} kg |
| 23a. | 21,000 Pa | 25c. | 1600 m/s |
| 23b. | Decreases to 61,000 Pa | 25d. | 744 kPa |
| 23c. | 5.8 km | | |

- 25e. 4.2×10^{20} or 0.0007 moles
 25g. 0.00785 m^3
 26a. 1.9 MW
 26b. 0.56 MW
 26c. 1.3 Mw
 27a. 54%
 27b. 240 kW
 27c. 890 kW
 27d. 590 kW
 27e. 630 kg
 28a. 98%
 28b. 4.0%
 28c. 12%
29. 14800 J
 30. 12,000 J
 31b. 720 K, 300 K, 600 K
 31c. isochoric; isobaric
 31d. C to A; B-C
 31e. 0.018 J
 32b. 300 K, 1200 K
 33a. 1753 J
 33b. -120 J
 33c. 80 J
 33d. 35 J
 33e. -100 J, 80 J, 80 J

Ch 20: Special and General Relativity

1. longer
 2. $\gamma = \infty$, the universe would be a dot
 3. 76.4 m, 76.4 m
 5. $\gamma = 1.002$
 6. $9.15 \times 10^7 \text{ m/s}$
 7. $2.6 \times 10^8 \text{ m/s}$
 8a. 0.659 km
 8b. 22.4
 8c. $4.92 \times 10^{-5} \text{ m/s}$
 8d. 14.7 km
9. 2900 m
 10. $1.34 \times 10^{-57} \text{ m}$
 11. 0.303 s
 12. $2.9 \times 10^{-30} \text{ kg}$, yes harder to accelerate
 13a. f
 13b. c
 14. $4.5 \times 10^{16} \text{ J}$; $1.8 \times 10^{13} \text{ softballs}$
 15a. $1.568 \times 10^{-13} \text{ J}$
 15b. $3.04 \times 10^6 \text{ J}$

Ch 21: Radioactivity and Nuclear Physics

- 6a. Substance A decays faster than B
 6b. Substance B because there is more material left to decay.
 7a. $^{219}_{88}\text{Ra} \rightarrow ^{215}_{86}\text{Rn} + ^4_2\text{He}$
 7b. $^{158}_{63}\text{Eu} \rightarrow ^{158}_{64}\text{Gd} + ^0_{-1}\text{e}^-$
 7c. $^{53}_{22}\text{Ti} \rightarrow ^{53}_{23}\text{Va} + ^0_{-1}\text{e}^-$
 7d. $^{211}_{83}\text{Bi} \rightarrow ^{207}_{81}\text{Tl} + ^4_2\text{He}$
 8a. 5×10^{24} atoms
 8b. Decay of a lot of atoms in a short period of time
 8c. 2.5×10^{24} atoms
 8d. $\frac{1}{2}$
 8e. 26.6 minutes
9. The one with the short half life, because half life is the rate of decay.
 10a. Substance B = 4.6 g and substance A = 0.035 g
 10b. substance B
 11. 1.2 g
 12. 125 g
 13. 0.46 minutes
 14. $t = 144,700$ years
 15. 0.0155 g
 16. 17 years
 17. 49,000 years

Ch 22: Standard Model of Particle Physics

1. strange
 2. Some type of meson
 3. Electron, photon, tau...
 4. Neutron, electron neutrino, Z^0
 5. Neutron, because it doesn't have electrical charge
 6. No, because it doesn't have electrical charge
7. Two anti-up quarks and an anti-down quark
 8. Lepton number, and energy/mass conservation
 9. Yes, W^+ , W^- , because they both have charge
 10. The weak force because it can interact with both quarks and leptons
 11. Yes; a,b,c,e; no; d,f

12. The standard model makes verifiable predictions, string theory makes few
- verifiable predictions.

Ch 23: Feynman Diagrams

1. Allowed: an electron and anti-electron(positron) annihilate to a photon then become an electron and anti-electron(positron) again.
2. Not allowed: electrons don't go backward though time, and charge is not conserved
3. Not allowed: lepton number is not conserved
- 4a. Allowed: two electrons exchange a photon
- 4b. Not allowed: neutrinos do not have charge and therefore cannot exchange a photon.
- 5a. Allowed: an electron and an up quark exchange a photon
- 5b. Not allowed: lepton number not conserved
6. Not allowed: quark number not conserved
7. Allowed: electron neutrino annihilates with a positron becomes a W^+ then splits to muon and muon neutrino.
8. Allowed: up quark annihilates with anti-up quark becomes Z^0 , then becomes a strange quark and anti-strange quark
9. Not allowed: charge not conserved
10. Allowed: this is a very rare interaction
11. Not allowed: electrons don't interact with gluons
12. Not allowed: neutrinos don't interact with photons
13. Allowed: the electron and the positron are exchanging virtual electron/positron pairs
14. Allowed: this is beta decay, a down quark splits into an up quark an electron and an electron neutrino via a W^- particle.
15. Allowed: a muon splits into an muon neutrino, an electron and an electron neutrino via a W^- particle.

Ch 24: Quantum Mechanics

- 1a. $6.75 \times 10^{-26} \text{ J}$, $2.253 \times 10^{-34} \text{ kgm/s}$
- 1b. $5.96 \times 10^{-20} \text{ J}$, $1.99 \times 10^{-28} \text{ kgm/s}$
- 1c. $4.90 \times 10^{-28} \text{ J}$, $1.63 \times 10^{-36} \text{ kgm/s}$
- 2a. 1.94 eV , $1.04 \times 10^{-27} \text{ kgm/s}$
- 2b. 12.7 eV , $6.76 \times 10^{-27} \text{ kgm/s}$
- 2c. 5.00 eV , $2.67 \times 10^{-21} \text{ kgm/s}$
- 3a. $.0827 \text{ nm}$
- 3b. $4.59 \times 10^{-4} \text{ nm}$
- 3c. 730 nm
4. $1.03 \times 10^{-20} \text{ m}$
- 5a. 36 nm
- 5b. no
- 5c. 380 nm , 73 nm , 36 nm , 92 nm , 39 nm , 74 nm
6. $.80 \text{ V}$
7. 480 nm
- 8a. $.124 \text{ nm}$
- 8b. $.00120 \text{ nm}$
9. $24,600 \text{ m/s}$
10. $1.84 \times 10^8 \text{ m/s}$
- 11a. $.491 \text{ m/s}$
- 11b. $3.14 \times 10^7 \text{ J}$
- 11c. 64 Mw
- 11d. 1.55 pm
12. 3.27 ev
- 14b. 15
- 14c. 182 nm , 188 nm , 206 nm , 230 nm
15. -10.3 ev , -3.82 eV , -2.29 eV , -1.83 eV
- 16a. $4.19 \times 10^7 \text{ m/s}$
- 16b. $1.70 \times 10^{-11} \text{ m}$
- 16c. 1.95°
- 16d. $.068 \text{ m}$
- 17a. 1.89 V
- 17b. 1.60 A
- 17c. 1.25Ω
- 18a. $4.40 \times 10^{-24} \text{ kgm/s}$
- 18b. $1.17 \times 10^{-24} \text{ kgm/s}$
- 18c. $3.23 \times 10^{-24} \text{ kgm/s}$
- 18d. $3.76 \times 10^7 \text{ m/s}$
- 19a. $1.1365 \times 10^{-22} \text{ kgm/s}$
- 19b. 5.860 pm
- 19c. $^{242}\text{Cu} \rightarrow {}^4\text{He} + {}^{238}\text{Pu}$
- 19d. 238.0497 amu
- 19e. 17.7 cm
- 19f. $-y$
- 19g. $+y$, 34.2 N/C
- 20.
21. $10^{-11} \text{ m} = 0.10 \text{ Angstrom}$
22. $1.7 \times 10^{-34} \text{ m}$