

UTTG-01-09

TCC-013-09

Living With Infinities

Steven Weinberg*

*Theory Group, Department of Physics, and Texas Cosmology Center,
University of Texas, Austin, TX, 78712*

Abstract

This is the written version of a talk given in memory of Gunnar Källén, at the Departments of Theoretical Physics, Physics, and Astronomy of Lund University on February 13, 2009. It will be published in a collection of the papers of Gunnar Källén, edited by C. Jarlskog and A. C. T. Wu. I discuss some of Källén's work, especially regarding the problem of infinities in quantum field theory, and recount my own interactions with him. In addition, I describe for non-specialists the current status of the problem, and present my personal view on how it may be resolved in the future.

*Electronic address: weinberg@physics.utexas.edu

I owe a great debt of gratitude to Gunnar Källén. In the summer of 1954, having just finished my undergraduate studies at Cornell, I arrived at the Bohr Institute in Copenhagen, where Källén was a member of the Theoretical Study Division of CERN, which had not yet moved to Geneva. Richard Dalitz had advised me to go to Copenhagen partly because of the presence there of CERN. But my real reason for coming to Copenhagen with my wife was that we had just married, and thought that we could have a romantic year abroad before we returned to the U.S. for me to enter graduate school. I brought with me a bag of physics books to read, but I did not imagine that I could start original research. You see, I had the idea that before I started research on any topic, I first had to know everything that had been done in that area, and I knew that I was far from knowing everything about anything.

It wasn't long before people at the Institute let me know that everyone there was expected to be working on some sort of research. David Frisch, a visiting American nuclear physicist, kindly suggested that I do something on nuclear alpha decay, but nothing came of it.

Early in 1955 I heard that a young theorist named Källén was doing interesting things in quantum field theory, so I knocked on his office door, and asked him to suggest a research problem. As it happened, Källén did have a problem to suggest. A year earlier, Tsung-Dao Lee at Columbia had invented a clever field-theoretic model that could be solved exactly.¹ The model had some peculiarities, which I'll come back to. These problems did not at first seem fatal to Lee, but Källén joined with the great Wolfgang Pauli to show that scattering processes in the Lee model violate the principle of unitarity — that is, the sum of the probabilities for all the things that can happen when two particles collide did not always add up to 100%.² Now Källén wanted me to see if there were other things wrong with the Lee model.

With a great deal of patient help from Källén, I was able to show that there were states in the Lee model whose energies were complex — that is, not ordinary real numbers. I finished the work on the Danish freighter that took my wife and me back to the U.S., and soon after I started graduate school at Princeton I had published the work as my first research paper.³

¹T.D. Lee, Phys. Rev. 95, 1329 (1954).

²G. Källén and W. Pauli, Dan. Mat. Fys. Medd. 30, no. 7 (1955).

³S. Weinberg, Phys. Rev. 102, 285 (1956).

This was a pretty unimportant paper (I recently checked, and found that it has been cited just nine times in 53 years), but it was a big thing for me — I started to feel like a physicist, not a student.

Incidentally, Källén's kindness to me went beyond starting me in research. He and his wife had my wife and me to their house for dinner, and going to the bathroom there, I learned something about Källén that probably most of you don't know — he had hand towels embroidered with the Dirac equation. Mrs. Källén told me that they were a present from Pauli. Källén also introduced me to Pauli, but I didn't get any towels.

Even though I had benefited so much from Källén's suggestion of a research problem, I felt that there was something odd about it. Lee was then not a well-known theorist — his great work with Yang on parity violation and weak interactions was a few years in the future. Also, the Lee model was not intended to be a serious model of real particles. So why did Källén take the trouble to shoot it down, even to the extent of enlisting the collaboration of his friend Pauli? The explanation, which I understood only much later, has to do with a long-standing controversy about the future of quantum field theory, in which Källén was playing an important part.

The controversy concerned the significance of infinities in quantum field theory. The problem of infinities was anticipated in the first papers on quantum field theory by Heisenberg and Pauli,⁴ and then in 1930 infinite energy shifts were found in calculations of the effects of emitting and reabsorbing photons by free or bound electrons, by Waller⁵ and Oppenheimer.⁶ In both cases you have to integrate over the momenta of the photons, and the integrals diverge. During the 1930s it was widely believed that these infinities signified a breakdown of quantum electrodynamics at energies of the order of 100 MeV. This changed after the war, when new techniques of calculation were developed that manifestly preserved the principles of special relativity at every step, and it was recognized that the infinities could be absorbed into a redefinition, called a *renormalization*, of physical constants like the charge and mass of the electron.⁷ Dyson was able to show (with some technicalities

⁴W. Heisenberg and W. Pauli, Z. f. Physik 56, 1 (1929); 59, 168 (1930).

⁵I. Waller, Z. f. Physik 59, 168 (1930); 61, 721, 837 (1930); 62, 673 (1930)

⁶J. R. Oppenheimer, Phys. Rev. 35, 461 (1930).

⁷See articles by Bethe, Dyson, Feynman, Kramers, Lamb & Retherford, Schwinger, Tomonaga, and Weisskopf reprinted in *Quantum Electrodynamics*, ed. J. Schwinger (Dover Publications, Inc., New York, 1958).

cleared up later by Salam⁸ and me⁹) that in quantum electrodynamics and a limited class of other theories, the renormalization of a finite number of physical parameters would actually remove infinities in every order of perturbation theory — that is, in every term when we write any physical observable as an expansion in powers of the charge of the electron, or powers of similar parameters in other theories. Theories in which infinities are removed in this way are known as *renormalizable*. They can be recognized by the property that in renormalizable theories, in natural units in which Planck’s constant and the speed of light are unity, all of the constants multiplying terms in the Lagrangian are just pure numbers, like the charge of the electron, or have the units of positive powers of energy, like particle masses, but not negative powers of energy.¹⁰

The great success of calculations in quantum electrodynamics using the renormalization idea generated a new enthusiasm for quantum electrodynamics. After this change of mood, probably most theorists simply didn’t worry about having to deal with infinite renormalizations. Some theorists thought that these infinities were just a consequence of having expanded in powers of the electric charge of the electron, and that not only observables but even quantities like the “bare” electron charge (the charge appearing in the field equations of quantum electrodynamics) would be found to be finite when we learned how to calculate without perturbation theory. But at least two leading theorists had their doubts about this, and thought that the appearance of infinite renormalizations in perturbation theory was a symptom of a deeper problem, a problem not with perturbation theory but with quantum field theory itself. They were Lev Landau, and Gunnar Källén.

Källén’s first step in exploring this problem was in an important 1952

⁸A. Salam, Phys. Rev. 82, 217 (1951).

⁹S. Weinberg, Phys. Rev. 118, 838 (1959).

¹⁰The units of these constants of course depend on the units we assign to the field operators. In using this criterion for renormalizability, it is essential to use units for any field operator related to the asymptotic behaviour of its propagator; if the propagator goes like k^n for large four-momentum k , then the field must be assigned the units of energy to the power $n/2 + 2$. In particular, because of $k^\mu k^\nu / (k^2 + m^2)$ terms in the propagator of a massive vector field, for these purposes the field must be given the unconventional units of energy to the power $+2$, and any interaction of the field would be non-renormalizable, unless the field is coupled only to conserved currents for which the terms in the propagator proportional to $k^\mu k^\nu$ may be dropped.

paper,¹¹ in which he showed how to define quantities like the bare charge of the electron without the use of perturbation theory. To avoid the complications that arise from the vector nature of the electromagnetic field, I'll describe the essential points here using the easier example of a real scalar field $\varphi(x)$, studied a little later by Lehmann.¹² The quantity $-i\Delta'(p)$ known as the propagator, that in perturbation theory would be given by the sum of all Feynman diagrams with two external lines, carrying four-momenta p^μ and $-p^\mu$, can be defined without the use of perturbation theory by

$$\langle 0|T\{\varphi(x), \varphi(0)\}|0\rangle = -i \int \frac{d^4p}{(2\pi)^4} \Delta'(p) e^{ip \cdot x}, \quad (1)$$

where $|0\rangle$ is the physical vacuum state, and T denotes a time-ordered product, with $\varphi(x)$ to the left or right of $\varphi(0)$ according as the time x^0 is positive or negative. By inserting a complete set of states between the fields in the time-ordered product, one finds what has come to be called the Källén–Lehmann representation

$$\Delta'(p) = \frac{|N|^2}{p^2 + m^2} + \int \frac{\sigma(\mu) d\mu}{p^2 + \mu^2}, \quad (2)$$

where $\sigma(\mu^2) \geq 0$ is given by a sum over multiparticle states with total energy-momentum vector P^λ satisfying $-P^2 = \mu^2$, and N is defined by the matrix element of $\varphi(x)$ between the vacuum and a one-particle state of physical mass m and three-momentum \mathbf{k} :

$$\langle 0|\varphi(x)|\mathbf{k}\rangle = \frac{N e^{ik \cdot x}}{(2\pi)^{3/2} \sqrt{2k^0}}, \quad (3)$$

with $k^0 \equiv \sqrt{\mathbf{k}^2 + m^2}$. If $\varphi(x)$ is the “unrenormalized” field that appears in the quadratic part of the Lagrangian without any extra factors, then it satisfies the canonical commutation relation

$$[\dot{\varphi}(\mathbf{x}, t), \varphi(\mathbf{y}, t)] = -i\delta^3(\mathbf{x} - \mathbf{y}). \quad (4)$$

By taking the time derivative of Eq. (1) and then setting the time x^0 equal to zero and using the commutation relation (4), one obtains the sum rule

$$1 = |N|^2 + \int \sigma(\mu) d\mu. \quad (5)$$

¹¹G. Källén, *Helv. Phys. Acta* 25, 417 (1952).

¹²H. Lehman, *Nuovo Cimento* XI, 342 (1954).

One immediate consequence is that, since $|N|^2$ is necessarily positive, Eq. (5) gives an upper limit on the coupling of the field φ to multiparticle states

$$\int \sigma(\mu) d\mu \leq 1 . \quad (6)$$

I'll mention in passing that this upper limit is reached in the case $N = 0$, which only applies if $\varphi(x)$ does not appear in the Lagrangian at all — that is, if the particle in question is not elementary. Thus, in a sense, composite particles are coupled to their constituents more strongly than any possible elementary particle.

This kind of sum rule has proved very valuable in theoretical physics. For instance, if instead of a pair of scalar fields in Eq. (1) we consider pairs of conserved symmetry currents, then by using methods similar to Källén's, one gets what are called a spectral function sum rules,¹³ which have had useful applications, for instance in calculating the decays of vector mesons into electron-positron pairs.

What chiefly concerned Källén was the application of these methods to quantum electrodynamics. In his 1952 paper, Källén derived a sum rule like (5) for the electromagnetic field, with $Z_3 \equiv |N_\gamma|^2$ in place of $|N|^2$, where N_γ is the renormalization constant for the electromagnetic field. As in the scalar field theory, this sum rule (and the definition of Z_3 as an absolute value squared) shows that

$$0 \leq Z_3 < 1 . \quad (7)$$

This is especially important in electrodynamics, because Z_3 appears in the relation between the bare electronic charge e_B that appears in the field equations, and the physical charge e of the electron:

$$e^2 = Z_3 e_B^2 . \quad (8)$$

The fact that e^2 is less than e_B^2 has a well-known interpretation: it is due to the shielding of the bare charge by virtual positrons, which are pulled out of the vacuum along with virtual electrons, and unlike the virtual electrons are attracted to the real electron whose charge is being measured.

Now, in lowest order perturbation theory, we have

$$Z_3 = 1 - \frac{e^2}{6\pi^2} \ln \left(\frac{\Lambda}{m_e} \right) , \quad (9)$$

¹³S. Weinberg, Phys. Rev. Lett. 18, 507 (1967).

where Λ is an ultraviolet cut-off, put in as a limit on the energies of the virtual photons. This is all very well if we take Λ as a reasonable multiple of the electron mass m_e , but if the cut-off is taken greater than $m_e \exp(6\pi^2/e^2) \approx 10^{280} m_e$ (which is more than the total mass of the observable universe) then we are in trouble: In this case Eq. (9) gives Z_3 negative, contradicting the inequality (7). As Landau pointed out,¹⁴ this ridiculously large energy becomes much smaller if we take into account the fact that there are several species of charged elementary particles; for instance, if there are ν species of spin one-half particles with the same charge as the electron, then the factor 10^{280} is replaced with $10^{280/\nu}$. So if ν is, say, 10 or 20, the problem with the sign of Z_3 would set in at energies much closer to those with which we usually have to deal. But this is just lowest order perturbation theory — to see if there is really any problem, it is necessary to go beyond perturbation theory.

To explore this issue, Källén set out to see if the integral appearing in $1 - Z_3$, and not just its expansion in powers of e^2 , actually diverges in the absence of a cut-off. Of course, he could not evaluate the integral exactly, but since every kind of multiparticle state makes a positive contribution to the integrand, he could concentrate on the contribution of the simplest states, consisting of just an electron and a positron — if the integral of this contribution diverges, then the whole integral diverges. In evaluating this contribution, he had to assume that all renormalizations including the renormalization of the electron mass and field were finite. With this assumption, and some tricky interchanges of integrations, he found that the integral for $1 - Z_3$ does diverge. In this way, he reached his famous conclusion that at least one of the renormalization constants in quantum electrodynamics has to be infinite.¹⁵

Not everyone was convinced. To quote the Källén memorial statement of Paul Urban in 1969,¹⁶ “Indeed, other authors are in doubt about his famous proof that at least one of the renormalization constants has to be infinite, but so far no definite answer to this question has been found.” It should be noted that at the end of his 1953 paper, Källén had explicitly disavowed any claim to mathematical rigor. As far as I know, this issue has never been settled. Of

¹⁴L. Landau, in *Niels Bohr and the Development of Physics* (Pergamon Press, New York, 1955): p. 52.

¹⁵G. Källén, *Dan. Mat. Fys. Medd.* 27, no. 12 (1953).

¹⁶P. Urban, *Acta Physica Austriaca*, Suppl. 6 (1969).

course, the important question was not whether some of the renormalization constants are infinite for infinite cut-off, but whether something happens at very high energies, such as $10^{280}m_e$, to prevent the cut-off in quantum electrodynamics from being taken to infinity. I don't know if Källén ever expressed an opinion about it, but I suspect that he thought that quantum electrodynamics does break down at very high energies, and that he wanted to be the one who proved it.

Which brings me back to the Lee model. This is a model with two heavy particles, V and N , and a lighter particle θ , all with zero spin. The only interactions in the theory are ones in which V converts to $N + \theta$, or vice versa. No antiparticles are included, and the recoil energies of the V and N are neglected, so the model is non-relativistic, though the energy ω of a θ of momentum \mathbf{p} is given by the relativistic formula $\omega = \sqrt{\mathbf{p}^2 + m_\theta^2}$. The model is exactly soluble in sectors with just one or two particles. For instance, to find the complete amplitude for $V \rightarrow N + \theta$, one can sum the graphs for

$$V \rightarrow N + \theta \rightarrow V \rightarrow N + \theta \rightarrow V \rightarrow \cdots \rightarrow N + \theta ,$$

which is just a geometric series. One finds that, if the physical and bare V -particle states are normalized so that

$$\langle V, \text{phys} | V, \text{phys} \rangle = \langle V, \text{bare} | V, \text{bare} \rangle = 1 , \quad (10)$$

then we have an exact sum rule resembling (5):

$$1 = |N|^2 + \frac{|g|^2}{4\pi^2} \int_0^\Lambda \frac{k^2 dk}{\omega^3} , \quad (11)$$

where

$$N \equiv \langle V, \text{bare} | V, \text{phys} \rangle \quad (12)$$

Here Λ is again an ultraviolet cut-off, and g is the renormalized coupling for this vertex, related to the bare coupling g_B by the exact formula $g = Ng_B$. For $\Lambda \gg m_\theta$, the integral in Eq. (11) grows as $\ln \Lambda$, so if $g \neq 0$ then Λ cannot be arbitrarily large without violating the condition that $|N|^2 \geq 0$. This is just like the problem encountered in lowest-order quantum electrodynamics, except that here there is no use of perturbation theory, and hence no hope that the difficulty will go away when perturbation theory is dispensed with.

Despite this difficulty, Lee found that his model with $\Lambda \rightarrow \infty$ gave sensible results for some simple problems, like the calculation of the energy of the V

particle. In their 1955 paper, Källén and Pauli confronted the difficulty that $|N|^2$ then comes out negative, and recognized that for very large Λ this was necessarily a theory with an indefinite metric — that is, it is necessary to take all states with odd numbers of bare V particles with negative norm, while all other states with definite numbers of bare particles have positive norm. In particular, in place of (10), we must take $\langle V, \text{bare} | V, \text{bare} \rangle = -1$, while calculations show that the physical V state has positive norm, so that we can still normalize it so that $\langle V, \text{phys} | V, \text{phys} \rangle = +1$. (There is also another energy eigenstate formed as a superposition of bare V and $N + \theta$ states, that has negative norm.) Then in place of (11), we have

$$1 = -|N|^2 + \frac{g^2}{4\pi^2} \int_0^\Lambda \frac{k^2 dk}{\omega^3}, \quad (13)$$

which gives no problem for large Λ . The device of an indefinite metric had already been introduced by Dirac,¹⁷ for reasons having nothing to do with infinities (Dirac was trying to find a physical interpretation of the negative energy solutions of the relativistic wave equations for bosons), and Pauli¹⁸ had noticed that if we can introduce suitable negative signs into sums over states, it should be possible to avoid infinities altogether. I think that what Källén and Pauli in 1955 disliked about the indefinite metric was not that it solved the problem of infinities, but that it did so too easily, without having to worry about what really happens at very high energies and short distances, and this is why they took the trouble to show that it did lead to unphysical results in the Lee model.

Experience has justified Källén and Pauli's distrust of the indefinite metric. This device continues to appear in theoretical physics, but only where there is some symmetry principle that cancels the negative probability for producing states with negative norm by the positive probability for producing other unphysical states, so that the total probability of producing physical states still adds up to 100%. Thus, in the Lorentz-invariant quantization of the electromagnetic field by Gupta and Bleuler,¹⁹ the state of a timelike photon has negative norm, but gauge invariance insures that the negative

¹⁷P. A. M. Dirac, Proc. Roy. Soc. A180, 1 (1942).

¹⁸W. Pauli, Rev. Mod. Phys. 15, 175 (1943).

¹⁹S. N. Gupta, Proc. Phys. Soc. 58, 681 (1950); K. Bleuler, Helv. Phys. Acta 28, 567 (1950).

probability for the production of these unphysical photons with timelike polarization is canceled by the positive probability for the production of other unphysical photons, with longitudinal polarization. A similar cancelation occurs in the Lorentz invariant quantization of string theories, where the symmetry is conformal symmetry on the two-dimensional worldsheet of the string. But it seems that without any such symmetry, as in the Lee model, the indefinite metric does not work.

I should say a word about where we stand today regarding the survival of quantum electrodynamics and other field theories in the limit of very high cut-off. The appropriate formalism for addressing this question is the renormalization group formalism presented by Wilson²⁰ in 1971. When we calculate the logarithmic derivative of the bare electron charge $e_{B\Lambda}$ with respect to the cut-off Λ at a fixed renormalized charge, then the result for $\Lambda \gg m_e$ can only depend on $e_{B\Lambda}$, since there is no relevant quantity with the units of energy with which Λ can be compared. That is, $e_{B\Lambda}$ satisfies a differential equation of the form

$$\Lambda \frac{de_{B\Lambda}}{d\Lambda} = \beta(e_{B\Lambda}) . \quad (14)$$

The whole question then reduces to the behavior of the function $\beta(e)$. If it is positive and increases fast enough so that $\int^\infty de/\beta(e)$ converges, then the cut-off in quantum electrodynamics cannot be extended to a value greater than a finite energy E_∞ , given by

$$E_\infty = \mu \exp \left(\int_{e_{B\mu}}^\infty \frac{de}{\beta(e)} \right) , \quad (15)$$

with μ arbitrary. On the basis of an approximation in which in each order of perturbation theory one keeps only terms with the maximum number of large logarithms, Landau concluded in ref. 14 that quantum electrodynamics does break down at very high energy. In effect, he was arguing on the basis of the lowest-order term, $\beta(e) \simeq e^3/12\pi^2$, for which $\int^\infty de/\beta(e)$ does converge.

No one today knows whether this is the case. It is equally possible that higher-order effects will make $\beta(e)$ increase more slowly or even decrease for very large e , in which case $\int^\infty de/\beta(e)$ will diverge and $e_{B\Lambda}$ just continue to grow smoothly with Λ . It is also possible that $\beta(e)$ drops to zero at some

²⁰K. G. Wilson, Phys. Rev. B4, 3174, 3184 (1971); Rev. Mod. Phys. 47, 773 (1975).

finite value e_* , in which case $e_{B\Lambda}$ approaches e_* as $\Lambda \rightarrow \infty$, though there are some arguments against this.²¹ Lattice calculations (in which spacetime is replaced by a lattice of separate points, providing an ultraviolet cut-off equal to the inverse lattice spacing) indicate that the beta function for a scalar field theory with interaction $g_B\varphi^4$ increases for large g_B fast enough so that $\int dg_B/\beta(g_B)$ converges and the theory therefore does not have a continuum limit for zero lattice spacing.²² And in the Lee model without an indefinite metric, Eq. (11) together with the relation $g_B = g/N$ gives

$$\beta(g_{B\Lambda}) \equiv \Lambda \frac{dg_{B\Lambda}}{d\Lambda} = \frac{g_{B\Lambda}^3}{8\pi^2}$$

for $\Lambda \gg m_\theta$, so $\int^\infty dg/\beta(g)$ converges, and as we have seen, the cut-off cannot be taken to infinity.

If limited to quantum electrodynamics, the problem of high energy behavior has become academic, since electromagnetism merges with the weak interactions at energies above 100 GeV, and we really should be asking about the high energy behavior of the $SU(2)$ and $U(1)$ couplings of the electroweak theory. Even that is somewhat academic, because gravitation becomes important at an energy of order 10^{19} GeV, well below the energy at which the $SU(2)$ and $U(1)$ couplings would become infinite. And there is no theory of gravitation that is renormalizable in the Dyson sense — the Newton constant appearing in General Relativity has the units of an energy to the power -2 .

Källén's concern with the problems of quantum field theory at very high energy did not keep him from appreciating the great success of quantum electrodynamics. In a contribution to the 1953 Kamerlingh Onnes Conference,²³ he remarked that “there is little doubt that the mathematical framework of quantum electrodynamics contains something which corresponds closely to physical reality.” He did practical calculations using perturbation theory in quantum electrodynamics, on problems such as the vacuum polarization in

²¹S. L. Adler, C. G. Callan, D. J. Gross, and R. Jackiw, Phys. Rev. D6, 2982 (1972); M. Baker and K. Johnson, Physica 96A, 120 (1979).

²²For a discussion and references, see J. Glimm and A. Jaffe, *Quantum Physics – A Functional Integral Point of View*, 2nd ed. (Springer-Verlag, New York, 1987), Sec. 21.6; R. Fernandez, J. Frölich, and A. D. Sokal, *Random Walks, Critical Phenomena, and Triality in Quantum Field Theory* (Springer-Verlag, Berlin, 1992), Chapter 15.

²³G. Källén, Physica XIX, 850 (1953).

fourth order²⁴ and the radiative corrections to decay processes.²⁵ He wrote a book about quantum electrodynamics,²⁶ leaving for the very end of the book his concern about the infinite value of renormalization constants.

Källén's interests were not limited to quantum electrodynamics. In 1954 he showed that the renormalizable meson theory with pseudoscalar coupling could not be used to account for both pion scattering and pion photoproduction, because different values of the pion-nucleon coupling constant are needed in the two cases.²⁷ Again, this result relied on lowest-order perturbation theory, so Källén acknowledged that it did not conclusively kill this meson theory. He remarked that "It would certainly be felt as a great relief by many theoretical physicists — among them the present author — if a definite argument against meson theory in its present form or a definite mathematical inconsistency in it could be found. This feeling together with wishful thinking must not tempt us to accept as conclusive evidence an argument that is still somewhat incomplete."

Of course, Källén was right in his distrust of this particular meson theory. A decade or so later the development of chiral Lagrangians showed that low energy pions are in fact well described by a theory with *pseudovector* coupling of single pions to nucleons, plus terms with two or more pions interacting with a nucleon at a single vertex, as dictated by a symmetry principle, chiral symmetry.²⁸ This theory is not renormalizable in the Dyson sense, but we have learned how to live with that. It is an effective field theory, which can be used to generate a series expansion for soft pion scattering amplitudes in powers of the pion energy. The Lagrangian for the theory contains every possible interaction that is allowed by the symmetries of the theory, but the non-renormalizable interactions whose coupling constants are negative powers of some characteristic energy (which is about 1 GeV in this theory) make a small contribution for pion energies that are much less than the characteristic energy. To any given order in pion energy, all infinities can be absorbed in the renormalization of a finite number of coupling parameters,

²⁴G. Källén and A. Sabry, Dan. Mat. Fys. Medd. 29, no. 7 (1955).

²⁵G. Källén, Nucl. Phys. B 1, 225 (1967).

²⁶G. Källén, *Quantum Electrodynamics*, transl. C. K. Iddings and M. Mizushima (Springer-Verlag, 1972).

²⁷G. Källén, Nuovo Cimento XII, 217 (1954).

²⁸For a discussion with references to the original literature, see Sec. 19.5 of S. Weinberg, *The Quantum Theory of Fields*, Vol. II (Cambridge Univ. Press, 1996.)

but we need more and more of these parameters to absorb infinities as we go to higher and higher powers of pion energy.

My own view is that all of the successful field theories of which we are so proud — electrodynamics, the electroweak theory, quantum chromodynamics, and even General Relativity — are in truth effective field theories, only with a much larger characteristic energy, something like the Planck energy, 10^{19} GeV. It is somewhat of an accident that the simplest versions of electrodynamics, the electroweak theory, and quantum chromodynamics are renormalizable in the Dyson sense, though it is very important from a practical point of view, because the renormalizable interactions dominate at ordinary accessible energies. An effect of one of the non-renormalizable terms has recently been detected: An interaction involving two lepton doublets and two scalar field doublets generates neutrino masses when the scalar fields acquire expectation values.²⁹

None of the renormalizable versions of these theories really describes nature at very high energy, where the non-renormalizable terms in the theory are not suppressed. From this point of view, the fact that General Relativity is not renormalizable in the Dyson sense is no more (or less) of a fundamental problem than the fact that non-renormalizable terms are present along with the usual renormalizable terms of the Standard Model. All of these theories lose their predictive power at a sufficiently high energy. The challenge for the future is to find the final underlying theory, to which the effective field theories of the standard model and General Relativity are low-energy approximations.

It is possible and perhaps likely that the ingredients of the underlying theory are not the quark and lepton and gauge boson fields of the Standard Model, but something quite different, such as a string theory. After all, as it has turned out, the ingredients of our modern theory of strong interactions are not the nucleon and pion fields of Källén's time, but quark and gluon fields, with an effective field theory of nucleon and pion fields useful only as a low-energy approximation.

But there is another possibility. The underlying theory may be an ordinary quantum field theory, including fields for gravitation and the ingredients of the Standard Model. Of course, it could not be renormalizable in the Dyson sense, so to deal with infinities every possible interaction allowed

²⁹S. Weinberg, *Phys. Rev. Lett.* 43, 1566 (1979).

by symmetry principles would have to be present, just as in effective field theories like the chiral theory of pions and nucleons. But it need not lose its predictive power at high energies, if the renormalized coupling constants $g_n(E)$ at a renormalization scale E approach a fixed point g_{n*} as $E \rightarrow \infty$.³⁰ This is known as “asymptotic safety.” For this to be possible, it is necessary that $\beta_n(g_*) = 0$, where $\beta_n(g(E)) \equiv E dg_n(E)/dE$. It is also necessary that the physical coupling constants $g_n(E)$ at any finite energy lie on a trajectory in coupling constant space that is attracted rather than repelled by this fixed point. There are reasons to expect that, even with an infinite number of coupling parameters, the surfaces spanned by such trajectories have finite dimensionality, so such a theory would involve just a finite number of free parameters, just as for ordinary renormalizable theories. The trouble, of course, is that there is no reason to expect the g_{n*} to be small, so that ordinary perturbation theory can’t be relied on for calculations in asymptotically safe theories. Other techniques such as dimensional continuation, $1/N$ expansions, and lattice quantization have provided increasing evidence that gravitation may be part of an asymptotically safe theory.³¹ So it is just possible that we may be closer to the final underlying theory than is usually thought.

Källén continued his interest in general elementary particle physics, and wrote a book about it, published in 1964.³² Arthur Wightman quoted a typical remark about this book: “That is the book on elementary particles that experimentalists find really helpful.” But Källén’s timing was unlucky – the development not only of chiral dynamics but of the electroweak theory were then just a few years in the future, and they were to put many of the problems he worried about in a new perspective.

It was a tragic loss not only to his friends and family but also to all theoretical physics that Källén died in an airplane accident just 40 years ago. For me, this was specially poignant, because he had been so kind to me in Copenhagen, and yet we had become estranged. Some time in 1957, just before I finished my graduate work, Källén visited Princeton, and left a note

³⁰S. Weinberg, in *General Relativity*, ed. S. W. Hawking and W. Israel (Cambridge University Press, 1979) 790.

³¹M. Reuter and F. Saueressig, 0708.1317; R. Percacci, in *Approaches to Quantum Gravity: Towards a New Understanding of Space, Time, and Matter*, ed. D. Oriti (Cambridge Univ. Press) [0709.3851]; D. F. Litim, 0810.3675; and earlier references cited therein.

³²G. Källén, *Elementary Particle Physics* (Addison-Wesley, Reading, MA. 1964).

in my mail box. Apparently he had seen a draft of my Ph. D. thesis, which was about the use of renormalization theory to deal with strong interaction effects in weak decay processes. His note seemed angry, and said that my work showed all the misconceptions about quantum field theory that were then common. Well, my thesis was no great accomplishment, but I didn't see why he was angry about it. Perhaps he was annoyed that I was following the common practice, of not worrying about the fact that the renormalization constants I encountered were infinite. Torsten Gustafson³³ has said of Källén that "Like Pauli he often expressed his opinion in a provocative fashion — especially to well-known physicists." I certainly was not a well-known physicist, but maybe Källén was paying me a compliment by treating me like one.

I did not meet Källén again after this, and I never replied to his note. I regret that very much, because I think that if I had replied we could have understood each other, and been friends again. Perhaps this talk can substitute for the reply to Källén that I should have made half a century ago.

I am grateful to C. Jarlskog and the Källén Lecture Committee for inviting me to Lund to give this talk, and to the Gunnar and Gunnel Källén Memorial Fund of the Royal Physiographic Society for sponsoring it. This material is based in part upon work supported by the National Science Foundation under Grant No. PHY-0455649 and with support from The Robert A. Welch Foundation, Grant No. F-0014

³³T. Gustafson, Nucl. Phys. A140, 1 (1970).