

*DUE: TUESDAY, JANUARY 20, 2015*

1. Show that for complex scalar fields,

$$\begin{aligned} \int \mathcal{D}\Phi^* \mathcal{D}\Phi \exp \left\{ i \int d^4x d^4y [\Phi^*(x) M(x, y) \Phi(y)] + i \int d^4x [J^*(x) \Phi(x) + \Phi^*(x) J(x)] \right\} \\ = \mathcal{N} \frac{1}{\det M} \exp \left\{ -i \int d^4x d^4y J^*(x) M^{-1}(x, y) J(y) \right\}, \end{aligned}$$

for some infinite constant  $\mathcal{N}$ . This is problem 14.1 on p. 283 of Schwartz.

2. (a) Derive the result:

$$\int d^4z \frac{\delta^2 W[J]}{\delta J(x) \delta J(z)} \frac{\delta^2 \Gamma[\Phi]}{\delta \Phi(z) \delta \Phi(y)} = -\delta^4(x - y),$$

and interpret diagrammatically. Here,  $W[J]$  is the generating functional for the connected Green functions and  $\Gamma[\Phi]$  is the generating functional for the one particle irreducible (1PI) Green functions.

(b) By taking one further functional derivative, show that  $\Gamma$  generates the amputated connected three-point function.

3. Consider the quantum field theory of a real scalar field governed by the Lagrangian,

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2 - \frac{\lambda}{4!} \phi^4.$$

(a) Evaluate the generating functional for the connected Green functions,  $W[J]$ , perturbatively, keeping all terms up to and including terms of  $\mathcal{O}(\lambda)$ .

(b) Using the result of part (a), compute the four-point connected Green function, take the appropriate Fourier transform, and verify the momentum space Feynman rule for the four-point scalar interaction obtained in class.

(c) Evaluate the classical field  $\phi_c(x)$  and the generating functional for the 1PI Green functions,  $\Gamma[\phi_c]$ , perturbatively, keeping all terms up to and including terms of  $\mathcal{O}(\lambda)$ . Then, repeat part (b) for the four-point 1PI Green function.

4. Consider a scalar field theory defined by the Lagrangian density

$$\mathcal{L} = \frac{1}{2} \partial^\mu \phi(x) \partial_\mu \phi(x) - V(\phi(x)), \quad (1)$$

and the corresponding equation of motion,

$$\square \phi(x) + V'(\phi) = 0,$$

where  $\square \equiv \partial^\mu \partial_\mu$  and  $V' \equiv dV/d\phi$ .

(a) Starting from eq. (14.122) on p. 276 of Schwartz, derive the equation of motion for the Green function  $\langle \Omega | T \{ \phi(x) \phi(y) \} | \Omega \rangle$ ,

$$\square_x \langle \Omega | T \{ \phi(x) \phi(y) \} | \Omega \rangle = -\langle \Omega | T \{ V'(\phi(x)) \phi(y) \} | \Omega \rangle - i \delta^4(x - y). \quad (2)$$

(b) Derive eq. (2) by the following technique. Start from the path integral definition of the generating functional,

$$Z[J] = \mathcal{N} \int \mathcal{D}\phi \exp \left\{ i \int d^4x [\mathcal{L} + J(x)\phi(x)] \right\}, \quad (3)$$

where  $\mathcal{N}$  is chosen such that  $Z[0] = 1$ . Perform a change of variables in the path integral,  $\phi(x) \rightarrow \phi(x) + \varepsilon(x)$ , where  $\varepsilon(x)$  is an arbitrary infinitesimal function of  $x$ . Noting that a change of variables<sup>1</sup> does not change the value of  $Z[J]$ , show that to first order in  $\varepsilon(x)$ ,

$$\int \mathcal{D}\phi \exp \left\{ i \int d^4x [\mathcal{L} + J(x)\phi(x)] \right\} \int d^4x \varepsilon(x) [-\square \phi - V'(\phi) + J(x)] = 0. \quad (4)$$

Since  $\varepsilon(x)$  is arbitrary, we may choose  $\varepsilon(x) = \epsilon \delta^4(x - y)$ , where  $\epsilon$  is an infinitesimal constant. With this choice for  $\varepsilon(x)$ , show that by taking the functional derivative of the eq. (4) with respect to  $J(x)$  and then setting  $J = 0$ , one can derive eq. (2).

*HINT:* What is the Jacobian corresponding to the change of variables,  $\phi(x) \rightarrow \phi(x) + \varepsilon(x)$ ?

5. Consider a field theory of a real pseudoscalar field coupled to a fermion field. The interaction Lagrangian is:

$$\mathcal{L}_{\text{int}} = -i\lambda \bar{\psi}(x) \gamma_5 \psi(x) \phi(x),$$

where  $\lambda$  is a real coupling constant (called the Yukawa coupling). Using functional techniques, derive the Feynman rule for the interaction vertex of this theory.

---

<sup>1</sup>Just as in the case of ordinary functional integration, a change of integration variables does not change the value of the functional integral.