

*DUE: TUESDAY, MAY 7, 2019*

**ALERT:** You should be ready with an initial choice for a term project topic on Monday April 29. Feel free to consult with me on possible choices. A short written proposal (one paragraph would suffice plus references) is due on Tuesday May 7.

1. The matrix group  $O(n)$  consists of real orthogonal  $n \times n$  matrices ( $n$  is a positive integer), and  $SO(n)$  consists of the subgroup of  $O(n)$  matrices with determinant equal to one.

(a) Show that  $SO(n)$  is a normal subgroup of  $O(n)$ .

(b) If  $n$  is odd, show that  $\mathbb{Z}_2 \cong \{\mathbf{I}_n, -\mathbf{I}_n\}$  is a normal subgroup of  $O(n)$ , where  $\mathbf{I}_n$  is the  $n \times n$  identity matrix. Prove that  $O(n)$  can be written as an internal direct product,  $O(n) \cong SO(n) \otimes \mathbb{Z}_2$ .

(c) Explain why the results of part (b) do not apply to the case of even  $n$ . Show that if  $n$  is even then  $O(n)$  can be written as a semidirect product,  $O(n) \cong SO(n) \rtimes \mathbb{Z}_2$ . Identify explicitly the subgroup of  $O(n)$  appearing in this semidirect product that is isomorphic to  $\mathbb{Z}_2$ .

2. A finite group  $G$  can be decomposed into conjugacy classes  $\mathcal{C}_k$ .

(a) Construct the set  $\mathcal{C}'_k \equiv g\mathcal{C}_k g^{-1}$ , which is obtained by replacing each element  $x \in \mathcal{C}_k$  by  $gxg^{-1}$ . Prove that  $\mathcal{C}'_k = \mathcal{C}_k$ .

(b) Suppose that  $D^{(i)}(g)$  is the  $i$ th irreducible (finite-dimensional) matrix representation of the finite group  $G$ . For a fixed class  $\mathcal{C}_k$ , prove that

$$\sum_{g \in \mathcal{C}_k} D_{j\ell}^{(i)}(g) = \frac{N_k}{n_i} \chi^{(i)}(\mathcal{C}_k) \delta_{j\ell}, \quad (1)$$

where  $n_i$  is the dimension of the  $i$ th irreducible representation of  $G$ ,  $N_k$  is the number of elements in the  $k$ th conjugacy class and  $\chi^{(i)}(\mathcal{C}_k)$  is the irreducible character corresponding to the  $k$ th conjugacy class.

*HINT:* Denoting the sum on the left hand side of eq. (1) by  $A_k$  and using the result of part (a), prove that  $D^{(i)}(g)A_k = A_k D^{(i)}(g)$  for all  $g \in G$ . Then use Schur's second lemma.

(c) Starting from the completeness result that is satisfied by the matrix elements of the irreducible matrix representations of  $G$  and using the result of part (b), derive the completeness relation for the irreducible characters,

$$\frac{N_k}{O(G)} \sum_i \chi^{(i)}(\mathcal{C}_k) [\chi^{(i)}(\mathcal{C}_\ell)]^* = \delta_{k\ell},$$

where  $O(G)$  is the order of the group  $G$  (i.e. the number of elements of  $G$ ), and the sum is taken over all inequivalent finite-dimensional irreducible representations.

(d) Using the orthogonality and the completeness relations satisfied by the irreducible characters, prove that the number of inequivalent irreducible representations of  $G$  is equal to the number of conjugacy classes.

3. Consider the transformations of the triangle that make up the dihedral group  $D_3$ . The elements of this group are  $D_3 = \{e, r, r^2, d, rd, r^2d\}$ , with the group multiplication law determined by the relations  $r^3 = e$ ,  $d^2 = e$  and  $dr = r^2d$ , where  $e$  is the identity element. In class, the following two-dimensional representation matrices for  $r, d \in D_n$  were given,

$$r = \begin{pmatrix} \cos(2\pi/n) & -\sin(2\pi/n) \\ \sin(2\pi/n) & \cos(2\pi/n) \end{pmatrix}, \quad d = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2)$$

Setting  $n = 3$ , one can construct a two-dimensional matrix representation of  $D_3$ .

(a) Consider the six-dimensional function space  $W$  consisting of polynomials of degree 2 in two real variables  $(x, y)$ :

$$f(x, y) = ax^2 + bxy + cy^2 + dx + ey + h, \quad (3)$$

where  $a, b, \dots, h$  are complex constants. We can view  $(a, b, \dots, h)$  as a six-dimensional vector that lives in a vector space which is isomorphic to  $W$ . If we perform a transformation of  $(x, y)$  under  $D_3$  according to the two-dimensional representation obtained from eq (2) with  $n = 3$ , then the polynomial  $f(x, y)$  given by eq. (3) transforms into another polynomial. That is, the vector  $(a, b, \dots, h)$  transforms under  $D_3$  according to a six-dimensional representation. Compute the  $6 \times 6$  matrices that represent the elements of  $D_3$ . Determine which irreducible representations of  $D_3$  are contained in this six-dimensional representation and their corresponding multiplicities.

(b) Identify the irreducible invariant subspaces of  $W$  under  $D_3$ . Check that your result is consistent with the results of part (a).

4. Consider the dihedral group  $D_4$  treated in problem 3 of Problem Set 1. The elements of this group are  $D_4 = \{e, r, r^2, r^3, d, rd, r^2d, r^3d\}$  with the group multiplication law determined by the relations  $r^4 = e$ ,  $d^2 = e$  and  $dr = r^3d$ , where  $e$  is the identity element.

(a) Write out the conjugacy class multiplication table.

(b) Determine explicitly the matrices of the regular representation.

(c) Using the two dimensional matrix representation given in eq. (2) with  $n = 4$ , verify that the group multiplication table of  $D_4$  is preserved. Prove that this representation is irreducible.

(d) Construct the character table for the irreducible representations of  $D_4$ .

5. Suppose that  $D$  is an irreducible  $n$ -dimensional representation of a finite group  $G$ , and  $D^{(1)}$  is a (nontrivial) one-dimensional representation of  $G$ . Prove that the direct product  $D \otimes D^{(1)}$  is an irreducible representation of  $G$ . [HINT: Show that for any  $g \in G$ ,  $|D^{(1)}(g)| = 1$ .]

6. (a) Display all the standard Young tableaux of the permutation group  $S_4$ . From this result, enumerate the inequivalent irreducible representations of  $S_4$  and specify their dimensions.

(b) Show that the normal subgroup  $\{e, (12)(34), (13)(24), (14)(23)\}$  of  $S_4$  is isomorphic to  $D_2$ . Using this result, prove that  $D_3 \cong S_4/D_2$ .

(c) Using the two-dimensional irreducible representation of  $D_3$  given in class and the result of part (b), construct a two-dimensional representation of  $S_4$  and determine its characters. Is the latter an *irreducible* representation of  $S_4$ ?

HINT: Show that given a normal subgroup  $N$  of a group  $G$  and a representation  $D^{G/N}$  of the factor group  $G/N$ , one can construct a representation  $D^G$  of the group  $G$  by defining  $D^G(g) \equiv D^{G/N}(gN)$  for all  $g \in G$ .

(d) Using the known one-dimensional representations of  $S_4$  and the results of parts (a) and (c), construct the character table for the group  $S_4$ . Determine any unknown entries in the character table by using the orthonormality and completeness relations for the irreducible characters. Using this technique, all entries of the character table can be uniquely determined up to a sign ambiguity in some of the entries.

(e) [EXTRA CREDIT] Resolve the sign ambiguity of part (d). For example, determine the matrix representative of the transposition (1 2) in the three-dimensional irreducible representation of  $S_4$ . By taking the trace of this matrix, complete the character table of  $S_4$ .

7. (a) Verify the following properties of the Pauli matrices  $\vec{\sigma} \equiv (\sigma_1, \sigma_2, \sigma_3)$ :

$$\begin{aligned} (i) \quad & \sigma_i \sigma_j = I \delta_{ij} + i \epsilon_{ijk} \sigma_k, \\ (ii) \quad & \sigma_2 \vec{\sigma} \sigma_2 = -\vec{\sigma}^*, \\ (iii) \quad & \exp(-i\theta \hat{\mathbf{n}} \cdot \vec{\sigma}/2) = I \cos(\theta/2) - i \hat{\mathbf{n}} \cdot \vec{\sigma} \sin(\theta/2), \end{aligned}$$

where  $I$  is the  $2 \times 2$  identity matrix.

(b) In the angle-and-axis parameterization of  $SO(3)$ , a rotation by an angle  $\theta$  about an axis that points along the unit vector  $\hat{\mathbf{n}}$  is represented by an  $SO(3)$  matrix given by  $R_{ij}(\hat{\mathbf{n}}, \theta) = \exp(-i\theta \hat{\mathbf{n}} \cdot \vec{\mathbf{J}})_{ij}$ , with  $(\hat{\mathbf{n}} \cdot \vec{\mathbf{J}})_{ij} \equiv -i \epsilon_{ijk} n_k$ . By convention, we assume that  $0 \leq \theta \leq \pi$ , and the axis  $\hat{\mathbf{n}}$  can point in any direction. Evaluate  $R_{ij}$  explicitly by summing the Taylor series of the exponential, and show that

$$R_{ij}(\hat{\mathbf{n}}, \theta) = n_i n_j + (\delta_{ij} - n_i n_j) \cos \theta - \epsilon_{ijk} n_k \sin \theta.$$

(c) Verify the formula:  $e^{-i\theta \hat{\mathbf{n}} \cdot \vec{\sigma}/2} \sigma_j e^{i\theta \hat{\mathbf{n}} \cdot \vec{\sigma}/2} = R_{ij}(\hat{\mathbf{n}}, \theta) \sigma_i$ .

(d) The set of matrices  $\exp(-i\theta \hat{\mathbf{n}} \cdot \vec{\sigma}/2)$  constitutes the defining representation of  $SU(2)$ . Prove that this representation is pseudoreal. [HINT: Property (ii) of part (a) is useful here.]