

Solution to the Equation of Motion for Forced Oscillations

The equation of motion for forced oscillations is given by Eq. (14-21) of Giancoli:

$$m \frac{dx^2}{dt^2} + b \frac{dx}{dt} + kx = F_0 \cos \omega t$$

We shall show that $x = A_0 \sin(\omega t + \phi_0)$ is a solution of by direct substitution.

$$x = A_0 \sin(\omega t + \phi_0) ; \frac{dx}{dt} = \omega A_0 \cos(\omega t + \phi_0) ; \frac{d^2x}{dt^2} = -\omega^2 A_0 \sin(\omega t + \phi_0)$$

$$m \frac{dx^2}{dt^2} + b \frac{dx}{dt} + kx = F_0 \cos \omega t \rightarrow$$

$$m[-\omega^2 A_0 \sin(\omega t + \phi_0)] + b[\omega A_0 \cos(\omega t + \phi_0)] + k[A_0 \sin(\omega t + \phi_0)] = F_0 \cos \omega t$$

Expanding the trigonometric functions [cf. page A-4 of Appendix A of Giancoli],

$$(kA_0 - m\omega^2 A_0)[\sin \omega t \cos \phi_0 + \cos \omega t \sin \phi_0] + b\omega A_0 [\cos \omega t \cos \phi_0 - \sin \omega t \sin \phi_0] = F_0 \cos \omega t$$

We now group the various terms by their time dependence.


$$\begin{aligned} & \left[(kA_0 - m\omega^2 A_0) \cos \phi_0 - b\omega A_0 \sin \phi_0 \right] \sin \omega t + \left[(kA_0 - m\omega^2 A_0) \sin \phi_0 + b\omega A_0 \cos \phi_0 \right] \cos \omega t \\ & = F_0 \cos \omega t \end{aligned}$$

The above equation must be valid for all *time*, which means that the coefficients of the functions of t must be the same on both sides of the equation. Since there is no $\sin \omega t$ on the right side of the equation, the coefficient of $\sin \omega t$ must be 0.

$$(kA_0 - m\omega^2 A_0) \cos \phi_0 - b\omega A_0 \sin \phi_0 = 0 \rightarrow$$

$$\frac{\sin \phi_0}{\cos \phi_0} = \frac{kA_0 - m\omega^2 A_0}{b\omega A_0} = \frac{k - m\omega^2}{b\omega} = \frac{m\omega_0^2 - m\omega^2}{b\omega} = \frac{\omega_0^2 - \omega^2}{\omega b/m} = \tan \phi_0 \rightarrow \boxed{\phi_0 = \tan^{-1} \frac{\omega_0^2 - \omega^2}{\omega b/m}}$$

Thus we see that Eq. 14-24 of Giancoli is necessary for $x = A_0 \sin(\omega t + \phi_0)$ to be the solution. This can be illustrated with the diagram shown below.



Finally, we equate the coefficients of $\cos \omega t$.

$$(kA_0 - m\omega^2 A_0) \sin \phi_0 + b\omega A_0 \cos \phi_0 = F_0 \rightarrow$$

$$A_0 \left[(k - m\omega^2) \frac{(\omega_0^2 - \omega^2)}{\sqrt{(\omega_0^2 - \omega^2)^2 + \frac{\omega^2 b^2}{m^2}}} + b\omega \frac{\frac{\omega b}{m}}{\sqrt{(\omega_0^2 - \omega^2)^2 + \frac{\omega^2 b^2}{m^2}}} \right] = F_0 \rightarrow$$

$$A_0 m \left[\frac{(\omega_0^2 - \omega^2)^2}{\sqrt{(\omega_0^2 - \omega^2)^2 + \frac{\omega^2 b^2}{m^2}}} + \frac{\frac{\omega^2 b^2}{m^2}}{\sqrt{(\omega_0^2 - \omega^2)^2 + \frac{\omega^2 b^2}{m^2}}} \right] = F_0 \rightarrow$$

$$\boxed{A_0 = \frac{F_0}{m \left[\sqrt{(\omega_0^2 - \omega^2)^2 + \frac{\omega^2 b^2}{m^2}} \right]}}$$

Thus we see that Eq. 14-23 of Giancoli is also necessary for $x = A_0 \sin(\omega t + \phi_0)$ to be the solution.