

DUE: MONDAY JUNE 11, 2012

FINAL EXAM ALERT: The final exam will be take place from 8–11 am on Wednesday June 13, 2010 in ISB 235 (unless a more convenient time can be found). The exam will cover the entire course. During the exam, you may consult Shankar and Baym (and any third quantum mechanics book of your choosing), your class notes, and any of the homework solutions and class handouts that are posted on the course website.

1. Consider a two-level system with $E_1 < E_2$. There is a time-dependent potential that connects the two levels as follows:

$$V_{11} = V_{22} = 0, \quad V_{12} = \gamma e^{i\omega t}, \quad V_{21} = \gamma e^{-i\omega t} \quad (\gamma \text{ real}),$$

where $V_{ij} = \langle i|V|j\rangle$. At time $t = 0$, it is known that only the lower level is populated—that is, $c_1(0) = 1$ and $c_2(0) = 0$. Note that a general state of the system can be expressed as a linear combination of eigenstates of the unperturbed Hamiltonian (in the Schrodinger picture):

$$|\psi(t)\rangle = \sum_{n=1}^2 c_n(t) e^{-iE_n t/\hbar} |n\rangle$$

(a) Starting with the time-dependent Schrodinger equation, derive the following differential equation for $c_k(t)$:

$$i\hbar \frac{dc_k}{dt} = \sum_{n=1}^2 V_{kn}(t) e^{i\omega_{kn}t} c_n, \quad (k = 1, 2), \quad (1)$$

where $V_{kn}(t) \equiv \langle k|V(t)|n\rangle$ and $\hbar\omega_{kn} \equiv E_k - E_n$. By solving the above system of differential equations *exactly*, find $|c_1(t)|^2$ and $|c_2(t)|^2$ for $t > 0$.

HINT: It is convenient to define new coefficients,

$$c'_1(t) \equiv e^{i(\omega_{21}-\omega)t/2} c_1(t), \quad c'_2(t) \equiv e^{-i(\omega_{21}-\omega)t/2} c_2(t).$$

Then, show that eq. (1) reduces to a matrix differential equation of the form

$$i\hbar \frac{d}{dt} \begin{pmatrix} c'_1(t) \\ c'_2(t) \end{pmatrix} = A \begin{pmatrix} c'_1(t) \\ c'_2(t) \end{pmatrix}, \quad (2)$$

where A is a *time-independent* 2×2 traceless hermitian matrix. Verify that the solution to eq. (2) is

$$\begin{pmatrix} c'_1(t) \\ c'_2(t) \end{pmatrix} = e^{-iAt/\hbar} \begin{pmatrix} c'_1(0) \\ c'_2(0) \end{pmatrix}.$$

By writing $A = \vec{a} \cdot \vec{\sigma}$ (where the vector \vec{a} is uniquely determined), it is straightforward to compute $e^{-iAt/\hbar}$ and complete part (a) of the problem.

(b) Do the same problem using time-dependent perturbation theory to lowest nonvanishing order. Compare the two approaches for small values of γ . Treat the following two cases separately: (i) ω very different from ω_{21} , and (ii) ω close to ω_{21} .

2. This problem provides a crude model for the photoelectric effect. Consider the hydrogen atom in its ground state (you may neglect the spins of the electron and proton). At time $t = 0$, the atom is placed in a high frequency uniform electric field that points in the z -direction,

$$\vec{\mathcal{E}}(t) = \mathcal{E}_0 \hat{z} \sin \omega t.$$

We wish to compute the transition probability per unit time that an electron is ejected into a solid angle lying between Ω and $\Omega + d\Omega$.

(a) Determine the minimum frequency, ω_0 , of the field necessary to ionize the atom.

(b) Using Fermi's golden rule for the transition rate at first-order in time-dependent perturbation theory, obtain an expression for the transition rate per unit solid angle as a function of the polar angle θ of the ejected electron (measured with respect to the direction of the electric field).

HINT: The matrix element that appears in Fermi's golden rule describes a transition of the negative-energy bound electron in its ground state to a positive-energy "free" electron. The wave function of the latter is actually quite complicated, since one cannot really neglect the effects of the long-range Coulomb potential. Nevertheless, you should simplify the computation by assuming the wave function of the ejected electron is a free-particle plane wave, with wave number vector \vec{k} . (Note that the direction of \vec{k} corresponds to that of the ejected electron).

(c) Integrate the result of part (b) over all solid angles to obtain the total ionization rate as a function of the frequency of the field. Determine the value of ω [in terms of ω_0 obtained in part (a)] for which the total ionization rate is maximal.

3. Consider the spontaneous emission of an $E1$ photon by an excited atom. The magnetic quantum numbers (m and m') of the initial and final atomic state are measured with respect to a fixed z -axis. Suppose the magnetic quantum number of the atom decreases by one unit.

(a) Compute the angular distribution of the emitted photon.

(b) Determine the polarization of the photon emitted in the z -direction.

(c) Verify that the result of part (b) is consistent with angular momentum conservation for the whole (atom plus photon) system.

HINT: The material on pp. 282–285 of Baym should be helpful.

4. Consider the elastic scattering of photons off electrons in atoms, assuming that the incident photon energies are large compared to the atomic binding energies. However, you should assume that the photon wavelength is still substantially larger than a typical atomic radius.

(a) Using the quantum theory of radiation, argue that the \vec{A} field operator must occur at least twice in the matrix element in order that there be a non-zero contribution in perturbation theory.

(b) Treating the quadratic $\vec{A} \cdot \vec{A}$ term in the interaction Hamiltonian to *first order* in perturbation theory, compute the differential cross-section in the dipole approximation. Show that:

$$\frac{d\sigma}{d\Omega} = r_0^2 |\vec{\epsilon}_\lambda \cdot \vec{\epsilon}_{\lambda'}^*|^2,$$

where $r_0 \equiv e^2/(mc^2)$ is the classical radius of the electron.

(c) Compute the total cross-section, assuming that the initial photon beam is unpolarized and the polarization of the final state photon is not measured.

5. Consider a non-interacting gas of N fermions that occupies a cubical box of volume V . Assume that V and N are macroscopically large. Let $\Psi_s(\vec{x})$ be the field operator that annihilates a fermion of spin orientation s at position \vec{x} . The particle number density operator is given by

$$n(\vec{x}) = \sum_s \Psi_s^\dagger(\vec{x}) \Psi_s(\vec{x}),$$

and the total number operator is $N = \int d^3x n(\vec{x})$. The Fourier transform of the number density operator is defined by

$$n_{\vec{q}} = \int d^3x n(\vec{x}) e^{-i\vec{q} \cdot \vec{x}}.$$

(a) Show that $n_{\vec{q}}$ can be expressed in terms of fermion creation and annihilation operators as follows:

$$n_{\vec{q}} = \sum_{\vec{k}} \sum_s a_{\vec{k},s}^\dagger a_{\vec{k}+\vec{q},s}.$$

(b) The *static structure function* for non-interacting fermions is defined as:

$$S_0(\vec{q}) \equiv \frac{1}{N} \langle \Phi_0 | n_{\vec{q}} n_{-\vec{q}} | \Phi_0 \rangle,$$

where $|\Phi_0\rangle$ is the N -particle ground state of the fermion gas (cf. Baym pp. 424–425). Evaluate $S_0(\vec{q})$ explicitly in the continuum limit.

HINT: Consider the cases $\vec{q} = 0$ and $\vec{q} \neq 0$ separately. In the former case, $S_0(\vec{q})$ can be obtained by inspection. The latter case results in an integral that can be evaluated by careful consideration of the limits of integration. One should find that $S_0(\vec{q})$ is a continuous function for $\vec{q} \neq 0$ but is not continuous at $\vec{q} = 0$.