RG-stable parameter relations of a scalar field theory in absence of a symmetry

Howard E. Haber seminar at SCIPP May 20, 2025

reference:

H.E. Haber and P.M. Ferreira, <u>arXiv:2502.11011</u> [hep-ph], Eur. Phys. J. C **85**, 541 (2025).



# <u>Outline</u>

- 1. Introduction-a folk theorem
- 2. Classifying the symmetries of the 2HDM
- 3. Parameter relation without a symmetry?
- 4. A toy model of two real scalar fields
- 5. Complexification and realification
- 6. Symmetries of the complexified theory
- 7. The origin of RG-stable parameter relations
- 8. Future directions

# Introduction-a folk theorem

In quantum field theory, generic tree-level parameter relations are not stable under renormalization group (RG) running.

If a tree-level parameter relation is the result of an unbroken symmetry, then the corresponding relation is RG-stable.

In the case of a softly-broken symmetry, the tree-level relations satisfied by dimensionless couplings are RG stable, although they receive *finite* radiative corrections.

Is the converse of the above statements true?

Folk theorem: the presence of an RG-stable parameter relation implies the existence of a symmetry.

A recent result, obtained by Ferreira, Grzadkowski, Ogreid, and Osland in the context of two-Higgs doublet model (2HDM), appears to violate this folk theorem.<sup>1</sup>

These authors attempted to resurrect the folk theorem by proposing a rather bizarre symmetry. But, is this a legitimate way to save the folk theorem?

Spoiler alert: the answer is no! Nevertheless, there is some truth to the folk theorem as I shall demonstrate in this talk.

<sup>&</sup>lt;sup>1</sup>P.M. Ferreira, B. Grzadkowski, O.M. Ogreid and P. Osland, Eur. Phys. J. C **84** (2024) 234, arXiv:2306.02410 [hep-ph].

## Symmetries of the 2HDM scalar potential

Consider the bosonic sector of the 2HDM:

$$\mathscr{L} = \mathscr{L}_{\mathrm{KE}} - \mathcal{V}(\Phi_1, \Phi_2),$$

where  $\mathscr{L}_{\text{KE}} \equiv (D^{\mu}\Phi_{a})^{\dagger}D_{\mu}\Phi_{a}$  (summed implicitly over a = 1, 2) is written in terms of the SU(2)×U(1)<sub>Y</sub> covariant derivative  $D_{\mu}$ , and the scalar potential is given by:

$$\begin{split} \mathcal{V} &= m_{11}^2 \Phi_1^{\dagger} \Phi_1 + m_{22}^2 \Phi_2^{\dagger} \Phi_2 - [m_{12}^2 \Phi_1^{\dagger} \Phi_2 + \text{h.c.}] \\ &+ \frac{1}{2} \lambda_1 (\Phi_1^{\dagger} \Phi_1)^2 + \frac{1}{2} \lambda_2 (\Phi_2^{\dagger} \Phi_2)^2 + \lambda_3 (\Phi_1^{\dagger} \Phi_1) (\Phi_2^{\dagger} \Phi_2) + \lambda_4 (\Phi_1^{\dagger} \Phi_2) (\Phi_2^{\dagger} \Phi_1) \\ &+ \left\{ \frac{1}{2} \lambda_5 (\Phi_1^{\dagger} \Phi_2)^2 + \left[ \lambda_6 (\Phi_1^{\dagger} \Phi_1) + \lambda_7 (\Phi_2^{\dagger} \Phi_2) \right] \Phi_1^{\dagger} \Phi_2 + \text{h.c.} \right\} \,, \end{split}$$

where  $m_{11}^2$ ,  $m_{22}^2$ , and  $\lambda_1, \dots, \lambda_4$  are real and  $m_{12}^2$ ,  $\lambda_5$ ,  $\lambda_6$  and  $\lambda_7$  are potentially complex parameters.

Parameter relations of the scalar potential can be the result of a symmetry that preserves the form of  $\mathscr{L}_{KE}$ . Two classes of symmetries are possible:

(1) Higgs flavor (HF) symmetries:  $\Phi_a \rightarrow S_{ab}\Phi_b$ .

(2) generalized CP symmetries (GCP):  $\Phi_a \to X_{ab} \Phi_b^*$ .

All possible inequivalent symmetries of the 2HDM scalar potential have been classified and the corresponding parameter relations elucidated.<sup>2</sup> We do not distinguish between different symmetries that yield the same parameter relations.

<sup>&</sup>lt;sup>2</sup>I.P. Ivanov, Phys. Lett. B **632** (2006) 360, hep-ph/0507132; Phys. Rev. D **75** (2007) 035001, hep-ph/0609018; P.M. Ferreira, H.E. Haber, and J.P. Silva, Phys. Rev. D **79** (2009) 116004, arXiv:0902.1537 [hep-ph]; P.M. Ferreira et al., Int. J. Mod. Phys. A **26** (2011) 769, arXiv:1010.0935 [hep-ph].

(1) HF symmetries [subgroups of U(2)]

$$\begin{split} \mathbb{Z}_2: \Phi_1 \to \Phi_1, \quad \Phi_2 \to -\Phi_2, \\ \mathrm{U}(1): \Phi_1 \to \Phi_1, \quad \Phi_2 \to e^{i\theta} \Phi_2, \quad 0 < \theta < 2\pi, \\ \mathrm{U}(2)/\mathrm{U}(1)_Y: \Phi_a \to S_{ab} \Phi_b, \quad \text{with } S \in \mathrm{U}(2)/\mathrm{U}(1)_Y. \end{split}$$

(2) GCP symmetries

$$\begin{aligned} \text{GCP1}: \ \Phi_1 \to \Phi_1^*, \quad \Phi_2 \to \Phi_2^*, \\ \text{GCP2}: \ \Phi_1 \to \Phi_2^*, \quad \Phi_2 \to -\Phi_1^*, \\ \text{GCP3}: \begin{cases} \Phi_1 \to \Phi_1^* \cos \theta + \Phi_2^* \sin \theta \\ \Phi_2 \to \Phi_2^* \cos \theta - \Phi_1^* \sin \theta \end{cases}, \quad 0 < \theta < \frac{1}{2}\pi. \end{aligned}$$

In the case of GCP3, any choice of  $0 < \theta < \frac{1}{2}\pi$  imposes the same conditions on the scalar potential parameters.

If we now impose the symmetries listed above in the scalar field basis  $\{\Phi_1, \Phi_2\}$ , we obtain the parameter relations listed below.

symmetry	$m_{22}^2$	$m_{12}^2$	$\lambda_2$	$\lambda_4$	${ m Re}\lambda_5$	${ m Im}\lambda_5$	$\lambda_6$	$\lambda_7$
$\mathbb{Z}_2$		0					0	0
U(1)		0			0	0	0	0
$U(2)/U(1)_Y$	$m_{11}^2$	0	$\lambda_1$	$\lambda_1 - \lambda_3$	0	0	0	0
GCP1		real				0	real	real
GCP2	$m_{11}^2$	0	$\lambda_1$					$-\lambda_6$
GCP3	$m_{11}^2$	0	$\lambda_1$		$\lambda_1 - \lambda_3 - \lambda_4$	0	0	0

Empty entries above correspond to a lack of constraints on the corresponding parameters.

What about other possible symmetries? For example,

 $\Pi_2: \Phi_1 \longleftrightarrow \Phi_2,$  $\Pi'_2: \Phi_1 \to \Phi_2, \quad \Phi_2 \to -\Phi_1,$  $GCP1': \Phi_1 \to \Phi_2^*, \quad \Phi_2 \to \Phi_1^*,$ 

symmetry	$m_{22}^2$	$m_{12}^2$	$\lambda_2$	$\operatorname{Re}\lambda_5$	${ m Im}\lambda_5$	$\lambda_6$	$\lambda_7$
$\Pi_2$	$m_{11}^2$	real	$\lambda_1$		0		$\lambda_6^*$
$\mathbb{Z}_2\otimes\Pi_2$	$m_{11}^2$	0	$\lambda_1$		0	0	0
$U(1) {\otimes} \Pi_2$	$m_{11}^2$	0	$\lambda_1$	0	0	0	0
$\Pi'_2$	$m_{11}^2$	pure imaginary	$\lambda_1$		0		$-\lambda_6^*$
U(1)'	$m_{11}^2$	pure imaginary	$\lambda_1$	$\lambda_1 - \lambda_3 - \lambda_4$	0	0	0
$U(1)^{\prime\prime}$	$m_{11}^2$	real	$\lambda_1$	$\lambda_3 + \lambda_4 - \lambda_1$	0	0	0
GCP1'	$m_{11}^2$		$\lambda_1$				$\lambda_6$
GCP3′	$m_{11}^2$	0	$\lambda_1$	$\lambda_3 + \lambda_4 - \lambda_1$	0	0	0

Taken from H.E. Haber and J.P. Silva, Phys. Rev. D 103 (2021) 115012, arXiv:2102.07136 [hep-ph].

By a change of the scalar field basis,  $\Phi_a \rightarrow U_{ab} \Phi_b$ ,<sup>3</sup> each of the symmetries above is equivalent to one of the six symmetries of the previous table (with its corresponding parameter relations).

For example, GCP1' is equivalent to GCP1 in another basis even though GCP1' (unlike GCP1) does not enforce reality conditions on the potentially complex parameters  $m_{12}^2$ ,  $\lambda_5$ ,  $\lambda_6$ , and  $\lambda_7$ .

<sup>&</sup>lt;sup>3</sup>Here,  $U \in U(2)$  is the most general transformation that preserves the gauge-kinetic energy terms.

## RG-stable parameter relations due to a symmetry

Consider the following one-loop beta functions (neglecting the contributions of the gauge and Yukawa couplings):

$$\begin{split} &16\pi^2\beta_{m_{11}^2} = 3\lambda_1m_{11}^2 + (2\lambda_3 + \lambda_4) m_{22}^2 - 3\left(\lambda_6^*m_{12}^2 + \lambda_6m_{12}^{2*}\right), \\ &16\pi^2\beta_{m_{22}^2} = (2\lambda_3 + \lambda_4) m_{11}^2 + 3\lambda_2m_{22}^2 - 3\left(\lambda_7^*m_{12}^2 + \lambda_7m_{12}^{2*}\right), \\ &16\pi^2\beta_{m_{12}^2} = -3\left(\lambda_6m_{11}^2 + \lambda_7m_{22}^2\right) + (\lambda_3 + 2\lambda_4) m_{12}^2 + 3\lambda_5m_{12}^{2*}, \\ &16\pi^2\beta_{\lambda_1} = 6\lambda_1^2 + 2\lambda_3^2 + 2\lambda_3\lambda_4 + \lambda_4^2 + |\lambda_5|^2 + 12|\lambda_6|^2, \\ &16\pi^2\beta_{\lambda_2} = 6\lambda_2^2 + 2\lambda_3^2 + 2\lambda_3\lambda_4 + \lambda_4^2 + |\lambda_5|^2 + 12|\lambda_7|^2, \\ &16\pi^2\beta_{\lambda_5} = (\lambda_1 + \lambda_2 + 4\lambda_3 + 6\lambda_4) \lambda_5 + 5\left(\lambda_6^2 + \lambda_7^2\right) + 2\lambda_6\lambda_7, \\ &16\pi^2\beta_{\lambda_6} = (6\lambda_1 + 3\lambda_3 + 4\lambda_4) \lambda_6 + (3\lambda_3 + 2\lambda_4) \lambda_7 + 5\lambda_5\lambda_6^* + \lambda_5\lambda_7^*, \\ &16\pi^2\beta_{\lambda_7} = (6\lambda_2 + 3\lambda_3 + 4\lambda_4) \lambda_7 + (3\lambda_3 + 2\lambda_4) \lambda_6 + 5\lambda_5\lambda_7^* + \lambda_5\lambda_6^*. \end{split}$$

Example 1:

Parameter relations:  $m_{11}^2 = m_{22}^2$ ;  $m_{12}^2 = 0$ ;  $\lambda_1 = \lambda_2$ ;  $\lambda_6 = -\lambda_7$ These relations are a consequence of a GCP2 symmetry. One can check that<sup>4</sup>

$$\begin{split} \beta_{m_{11}^2 - m_{22}^2} \big|_{\text{sym}} &\equiv \left[ \beta_{m_{11}^2} - \beta_{m_{22}^2} \right] \big|_{\text{sym}} = 0 \,, \\ \beta_{m_{12}^2} \big|_{\text{sym}} &\equiv 0 \,, \\ \beta_{\lambda_1 - \lambda_2} \big|_{\text{sym}} &\equiv \left[ \beta_{\lambda_1} - \beta_{\lambda_2} \right] \big|_{\text{sym}} = 0 \,, \\ \beta_{\lambda_6 + \lambda_7} \big|_{\text{sym}} &\equiv \left[ \beta_{\lambda_6} + \beta_{\lambda_7} \right] \big|_{\text{sym}} = 0 \,, \end{split}$$

Indeed, the parameter relations given above are RG-stable to all orders in perturbation theory.

<sup>&</sup>lt;sup>4</sup>The notation "sym" indicates that the parameter relations were used when evaluating the beta functions.

The GCP2 symmetry guarantees that the RG-stability persists when the gauge interactions are included.<sup>5</sup>

Moreover, the RG-stability of the tree-level parameter relations persists to all orders of perturbation theory.

Example 2:6

Parameter relations:  $m_{11}^2 = -m_{22}^2$ ;  $\lambda_1 = \lambda_2$ ;  $\lambda_6 = -\lambda_7$ 

$$\begin{split} \beta_{m_{11}^2 + m_{22}^2} \Big|_{\text{sym}} &\equiv \left[ \beta_{m_{11}^2} + \beta_{m_{22}^2} \right] \Big|_{\text{sym}} = 0 \\ \beta_{\lambda_1 - \lambda_2} \Big|_{\text{sym}} &\equiv \left[ \beta_{\lambda_1} - \beta_{\lambda_2} \right] \Big|_{\text{sym}} = 0 , \\ \beta_{\lambda_6 + \lambda_7} \Big|_{\text{sym}} &\equiv \left[ \beta_{\lambda_6} + \beta_{\lambda_7} \right] \Big|_{\text{sym}} = 0 . \end{split}$$

)

<sup>&</sup>lt;sup>5</sup>If the GCP2 symmetry could be extended to the Yukawa sector, then the RG-stability would also persist when the Yukawa interactions are included.

<sup>&</sup>lt;sup>6</sup>First considered by P.M. Ferreira, B. Grzadkowski, O.M. Ogreid, and P. Osland, Eur. Phys. J. C **84** (2024) 234, arXiv:2306.02410 [hep-ph].

The parameter relations,  $\lambda_1 = \lambda_2$ ;  $\lambda_6 = -\lambda_7$ , are RG-stable to all orders in perturbation theory since the GCP2 symmetry is softly broken by the squared-mass terms of the scalar potential.

The parameter relation,  $m_{11}^2 = -m_{22}^2$  is RG-stable at one-loop order, and persists at two-loop order (and beyond).<sup>7</sup>

However, no basis change exists such that  $m_{11}^2 = -m_{22}^2$  is transformed into one of the six symmetry parameter relations previously classified.

Has a possible symmetry of the 2HDM scalar potential been overlooked?

<sup>&</sup>lt;sup>7</sup>As shown by P.M. Ferreira et al., op. cit., the RG-stability holds to all orders in perturbation theory in the quartic scalar couplings and the gauge couplings. It has also been shown to hold at two-loop order in the Yukawa couplings (but no corresponding all-orders result has yet been obtained).

# GOOFy symmetry?

<u>Reference</u>: B. Grzadkowski, O.M. Ogreid, P. Osland, and P.M. Ferreira, op. cit.

Consider the following scalar field transformations:

$$\Phi_1 \to -\Phi_2^*, \quad \Phi_1^* \to \Phi_2, \quad \Phi_2 \to \Phi_1^*, \quad \Phi_2^* \to -\Phi_1.$$

This is peculiar since  $\Phi_1^*$  does not transform into the complex conjugate of the transformed  $\Phi_1$  (and similarly for  $\Phi_2$ ). Equivalently,

$$\Phi_1^{\dagger}\Phi_1 
ightarrow -\Phi_2^{\dagger}\Phi_2\,, \qquad \Phi_2^{\dagger}\Phi_2 
ightarrow -\Phi_1^{\dagger}\Phi_1,$$

whereas  $\Phi_1^{\dagger}\Phi_2$  and  $\Phi_2^{\dagger}\Phi_1$  are invariant. Imposing this "GOOFy symmetry" on the scalar potential yields the parameter relations,  $m_{11}^2 = -m_{22}^2$ ;  $\lambda_1 = \lambda_2$ ;  $\lambda_6 = -\lambda_7$ .

Unfortunately,  $\mathscr{L}_{KE}$  changes sign under  $\Phi_1^{\dagger}\Phi_1 \rightarrow -\Phi_2^{\dagger}\Phi_2$  and  $\Phi_2^{\dagger}\Phi_2 \rightarrow -\Phi_1^{\dagger}\Phi_1$ .

In order to restore the sign of  $\mathscr{L}_{\text{KE}}$ , the authors of the GOOFy paper advanced the radical proposal where the spacetime coordinates themselves also transform under the GOOFy symmetry via  $x_{\mu} \rightarrow i x_{\mu}$ .<sup>8</sup>

My personal conclusion is that the GOOFy symmetry is not a legitimate symmetry. In particular, it seems that there is no valid symmetry explanation for the RG-stable parameter relation  $m_{11}^2 = -m_{22}^2$ .

<sup>&</sup>lt;sup>8</sup>Equivalently, the covariant derivative must also transform as  $D_{\mu} \rightarrow iD_{\mu}$  (which implies that the gauge fields themselves must also similarly transform) in order that the kinetic energy terms of the scalar fields remain invariant.

## A toy model with one complex scalar field

Consider a theory of one complex scalar field  $\Phi$  with Lagrangian,

$$\mathscr{L} = \partial_{\mu} \Phi \partial^{\mu} \Phi^{*} - m_{1}^{2} \Phi^{*} \Phi - m_{2}^{2} \Phi^{2} - m_{2}^{2*} \Phi^{*2} - \lambda_{1} (\Phi^{*} \Phi)^{2}$$
$$-\lambda_{2} \Phi^{4} - \lambda_{2}^{*} \Phi^{*4} - (\lambda_{3} \Phi^{2} + \lambda_{3}^{*} \Phi^{*2}) \Phi^{*} \Phi ,$$

after imposing a discrete symmetry  $\Phi \to -\Phi$  to remove terms linear and cubic in the scalar fields. Next, impose the relations:

$$m_1^2 = \lambda_3 = 0 \,.$$

These parameter relations are RG-stable to all orders in perturbative theory:

$$\beta_{m_1^2}\big|_{\text{sym}} = 0, \qquad \beta_{\lambda_3}\big|_{\text{sym}} = 0,$$

where "sym" means that the  $\beta$ 's are evaluated at  $m_1^2 = \lambda_3 = 0$ .

The symmetry transformation  $\Phi \rightarrow i\Phi$  would set  $m_2^2 = \lambda_3 = 0$ . This symmetry is softly broken, which explains why the relation  $\lambda_3 = 0$  is RG-stable. But why is  $m_1^2 = 0$  RG-stable?

#### A GOOFy-like symmetry?

The "symmetry" transformation  $\Phi \to \Phi$ ;  $\Phi^* \to -\Phi^*$ , removes the terms  $m_1^2 \Phi^* \Phi + (\lambda_3 \Phi^2 + \lambda_3^* \Phi^{*2}) \Phi^* \Phi$ . That is,  $m_1^2 = \lambda_3 = 0$ .

Once again, the complex conjugate of the  $\Phi$  transformation is not equal to the  $\Phi^*$  transformation. Moreover, one must again restore the sign of the kinetic energy term  $\mathscr{L}_{\text{KE}} = \partial_{\mu} \Phi \partial^{\mu} \Phi^*$  by transforming  $x_{\mu} \to i x_{\mu}$ . That is, there seems to be no legitimate symmetry explanation for the RG-stability of  $m_1^2 = 0$ .

## Realification of a complex scalar field theory

One can always "realify" a complex scalar field theory by writing  $\Phi = (\varphi_1 + i\varphi_2)/\sqrt{2}$ . After imposing the GOOFy-like symmetry,

$$\mathcal{V} = \frac{1}{2}m_{11}^2 \left(\varphi_1^2 - \varphi_2^2\right) + m_{12}^2 \varphi_1 \varphi_2 + \frac{1}{24}\lambda_{1111} \left(\varphi_1^4 + \varphi_2^4\right) \\ + \frac{1}{4}\lambda_{1122} \left(\varphi_1 \varphi_2\right)^2 + \frac{1}{6}\lambda_{1112} \left(\varphi_1^2 - \varphi_2^2\right) \varphi_1 \varphi_2.$$

In particular, there are three parameter relations:<sup>9</sup>

$$m_{22}^2 = -m_{11}^2$$
  $\lambda_{1111} = \lambda_{2222}$ ,  $\lambda_{1112} = -\lambda_{1222}$ ,

which are equivalent to the previous relations,  $m_1^2 = \lambda_3 = 0$ .

<sup>&</sup>lt;sup>9</sup>The notation corresponds to  $\mathcal{V} = \frac{1}{2}m_{ij}^2\varphi_i\varphi_j + \frac{1}{24}\lambda_{ijk\ell}\varphi_i\varphi_j\varphi_k\varphi_\ell$ , with an implicit sum over repeated indices. The specific model above was first proposed in a talk by B. Grzadkowski at the at the 2024 Workshop on Multi-Higgs Models in Lisbon, Portugal.

The symmetry transformations  $\varphi_1 \rightarrow \varphi_2$ ;  $\varphi_2 \rightarrow -\varphi_1$  would set  $m_{11}^2 = m_{22}^2$ ;  $m_{12}^2 = 0$ ;  $\lambda_{1111} = \lambda_{2222}$ ;  $\lambda_{1112} = -\lambda_{1222}$ . This symmetry is softly broken, so that the quartic coupling relations are RG-stable. But why is  $m_{11}^2 = -m_{22}^2$  RG-stable?

The corresponding GOOFy-like "symmetry" transformations that yield  $m_{11}^2 = -m_{22}^2$ ;  $\lambda_{1111} = \lambda_{2222}$ ;  $\lambda_{1112} = -\lambda_{1222}$  are:<sup>10</sup>

 $\varphi_1 \to i\varphi_2 , \qquad \qquad \varphi_2 \to -i\varphi_1 .$ 

This is not a legitimate symmetry of a *real* scalar field theory.<sup>11</sup> But, the parameter relation  $m_{22}^2 = -m_{11}^2$  is RG-stable to all orders of perturbation theory. Indeed,

$$\beta_{m_{11}^2 + m_{22}^2} \big|_{\text{sym}} = \beta_{m_{11}^2} + \beta_{m_{22}^2} \big|_{\text{sym}} = 0 \,,$$

<sup>10</sup>These relations are similar to the 2HDM parameter relations imposed by the GOOFy symmetry. <sup>11</sup>Moreover, this "symmetry" transformation flips the sign of the kinetic energy terms.

#### Some technical details

The formulae for the one-loop and two-loop beta functions,  $\beta \equiv \beta^I + \beta^{II}$ , of a real scalar field theory are:

$$\beta_{m_{ij}^2}^I = m_{mn}^2 \lambda_{ijmn} ,$$
  
$$\beta_{\lambda_{ijk\ell}}^I = \frac{1}{8} \sum_{\text{perm}} \lambda_{ijmn} \lambda_{mnk\ell} = \lambda_{ijmn} \lambda_{mnk\ell} + \lambda_{ikmn} \lambda_{mnj\ell} + \lambda_{i\ell mn} \lambda_{mnjk} ,$$

with an implicit sum over the repeated indices, where  $\sum_{\text{perm}}$  denotes a sum over the permutations of the uncontracted indices, i, j, k, and  $\ell$ , and

$$\beta_{m_{ij}^2}^{II} = \frac{1}{12} \left( \lambda_{ik\ell m} \lambda_{nk\ell m} m_{nj}^2 + \lambda_{jk\ell m} \lambda_{nk\ell m} m_{ni}^2 \right) - 2m_{k\ell}^2 \lambda_{ikmn} \lambda_{j\ell mn} ,$$
  
$$\beta_{\lambda_{ijk\ell}}^{II} = \frac{1}{72} \sum_{\text{perm}} \lambda_{inpq} \lambda_{mnpq} \lambda_{mjk\ell} - \frac{1}{4} \sum_{\text{perm}} \lambda_{ijmn} \lambda_{kmpq} \lambda_{\ell npq} .$$

The  $\beta^{II}$  above each consist of the sum of two linearly independent combinations of tensor quantities. Each individual combination separately vanishes when the parameter relations (indicated by "sym") are applied.

Explaining the mysterious RG-stability: Complexification

The illegitimate symmetry

$$\varphi_1 \to i\varphi_2, \qquad \qquad \varphi_2 \to -i\varphi_1.$$

would have been a legitimate symmetry if the  $\varphi_i$  had been complex scalar fields.

Our proposal is to promote the two real fields  $\varphi_1$  and  $\varphi_2$  to complex fields  $\Phi_1$  and  $\Phi_2$ , and then impose the (now legitimate) symmetry transformations

$$\Phi_1 \to i \Phi_2 , \qquad \Phi_2 \to -i \Phi_1 .$$

We will additionally impose a CP symmetry,  $\Phi_i \rightarrow \Phi_i^*$ , to impose reality conditions on the parameters of the scalar potential.

The resulting parameter relations of the complexified theory are RG-stable due to these symmetries.

Moreover, we will show that the corresponding  $\beta$ -function relations of the complexified theory can be related to  $\beta$ -function relations obtained in the original real scalar field theory.

That is, the RG-stability of the parameter relations of the original theory can be attributed to symmetries of the complexified theory, thereby restoring the folk theorem.

#### The complexification recipe

- Promote the real scalar fields  $\varphi_i$  to complex fields  $\Phi_i$ .
- The complexified model is *defined* to employ a canonically normalized kinetic energy term,

$$\mathscr{L}_{\rm KE} = \partial^{\mu} \Phi_{\bar{a}}^* \partial_{\mu} \Phi_a \,,$$

- Impose the appropriate  $\mathbb{Z}_2$  symmetry to eliminate terms with an odd number of fields.
- Promote the GOOFy-like symmetries of the real scalar field theory to legitimate symmetries of the complexified theory.
- Impose a CP symmetry so that the scalar potential parameters are real.

The kinetic energy term,  $\mathscr{L}_{KE} = \partial^{\mu} \Phi_{\bar{a}}^* \partial_{\mu} \Phi_a$ , is invariant under a U(2) basis transformation,<sup>12</sup>

$$\Phi_a \to U_{a\bar{b}} \Phi_b , \qquad \Phi^*_{\bar{a}} \to \Phi^*_{\bar{b}} U^{\dagger}_{b\bar{a}} ,$$

where  $U_{b\bar{a}}^{\dagger}U_{a\bar{c}} = \delta_{b\bar{c}}$ . In the notation introduced above, the indices  $a, b, c \in \{1, 2\}$  and  $\bar{a}, \bar{b}, \bar{c} \in \{\bar{1}, \bar{2}\}$  run over the complex two dimensional flavor space of the scalar fields. The use of the unbarred/barred index notation is accompanied by the rule that there is an implicit sum over unbarred/barred index pairs.

<sup>&</sup>lt;sup>12</sup>In a different (and perhaps more common) index convention, unbarred indices are lower indices and barred indices are upper indices, with the rule that one sums over upper/lower index pairs. For notational reasons, we preferred the unbarred/barred index notation in this work.

#### Scalar potential of the complexified model

After removing terms with an odd number of fields, the most general renormalizable scalar potential is:

$$\mathcal{V}_{C} = M_{a\bar{b}}^{2} \Phi_{\bar{a}}^{*} \Phi_{b} + M_{\bar{a}\bar{b}}^{2} \Phi_{a} \Phi_{b} + M_{ab}^{2} \Phi_{\bar{a}}^{*} \Phi_{\bar{b}}^{*} + \Lambda_{ab\bar{c}\bar{d}} \Phi_{\bar{a}}^{*} \Phi_{\bar{b}}^{*} \Phi_{c} \Phi_{d}$$
$$+ \Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}} \Phi_{a} \Phi_{b} \Phi_{c} \Phi_{d} + \Lambda_{a\bar{b}\bar{c}\bar{d}} \Phi_{\bar{a}}^{*} \Phi_{b} \Phi_{c} \Phi_{d} + \Lambda_{abc\bar{d}} \Phi_{\bar{a}}^{*} \Phi_{\bar{b}}^{*} \Phi_{\bar{c}}^{*} \Phi_{d} + \Lambda_{abcd} \Phi_{\bar{a}}^{*} \Phi_{\bar{b}}^{*} \Phi_{\bar{c}}^{*} \Phi_{d}^{*}$$

In this notation,  $M_{ab}^2$  and  $M_{a\bar{b}}^2$  are independent tensors (despite the use of the same symbol  $M^2$ ). Likewise,  $\Lambda_{abcd}$ ,  $\Lambda_{abc\bar{d}}$ , and  $\Lambda_{ab\bar{c}\bar{d}}$  are independent tensors (despite the use of the same symbol  $\Lambda$ ).

Hermiticity and permutation symmetry imply

$$M_{\bar{a}\bar{b}}^{2} = M_{\bar{b}\bar{a}}^{2}, \qquad M_{ab}^{2} = M_{ba}^{2}, \qquad \Lambda_{ab\bar{c}\bar{d}} = \Lambda_{ba\bar{c}\bar{d}} = \Lambda_{ab\bar{d}\bar{c}} = \Lambda_{ba\bar{d}\bar{c}},$$
$$M_{a\bar{b}\bar{c}}^{2} = \left[M_{b\bar{a}}^{2}\right]^{*}, \qquad \Lambda_{ab\bar{c}\bar{d}} = \left[\Lambda_{cd\bar{a}\bar{b}}\right]^{*},$$
$$M_{\bar{a}\bar{b}}^{2} = \left[M_{ab}^{2}\right]^{*}, \qquad \Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}} = \left[\Lambda_{abcd}\right]^{*}, \qquad \Lambda_{d\bar{a}\bar{b}\bar{c}} = \left[\Lambda_{abc\bar{d}}\right]^{*}.$$

In particular,  $M_{1\bar{1}}^2$ ,  $M_{2\bar{2}}^2$ ,  $\Lambda_{11\bar{1}\bar{1}}$ ,  $\Lambda_{22\bar{2}\bar{2}}$ , and  $\Lambda_{12\bar{1}\bar{2}} = \Lambda_{21\bar{2}\bar{1}}$  are real parameters.

Imposing the symmetry on the complexified model

If  $\mathcal{V}_C$  is invariant under  $\Phi_1 \rightarrow i \Phi_2$  and  $\Phi_2 \rightarrow -i \Phi_1$ , then the following parameter relations are obtained:

$$\begin{split} M_{1\bar{1}}^2 &= M_{2\bar{2}}^2 & M_{1\bar{2}} = 0, & M_{1\bar{1}}^2 = -M_{22}^2, \\ \Lambda_{1111} &= \Lambda_{2222}, & \Lambda_{1112} = -\Lambda_{1222}, \\ \Lambda_{111\bar{1}} &= -\Lambda_{222\bar{2}}, & \Lambda_{112\bar{1}} = \Lambda_{122\bar{2}}, & \Lambda_{112\bar{2}} = -\Lambda_{122\bar{1}}, & \Lambda_{111\bar{2}} = \Lambda_{222\bar{1}}, \\ \Lambda_{11\bar{1}\bar{1}} &= \Lambda_{22\bar{2}\bar{2}}, & \Lambda_{11\bar{1}\bar{2}} = -\Lambda_{12\bar{2}\bar{2}}, & \Lambda_{11\bar{2}\bar{2}} = \Lambda_{11\bar{2}\bar{2}}, \end{split}$$

After imposing a CP symmetry, which renders all scalar potential parameters real, were are left with three real squared mass terms and eleven real quartic couplings. The resulting scalar potential is depends on three real squared mass terms and eleven real quartic couplings:

$$\begin{split} M_{ab}^2 &= M_{\bar{a}\bar{b}}^2 \ni \{\bar{M}^2, M_{12}^2\}\,,\\ M_{a\bar{b}}^2 &= M_{b\bar{a}}^2 \ni \{M^2\}\,,\\ \Lambda_{ab\bar{c}\bar{d}} &= \Lambda_{cd\bar{a}\bar{b}} \ni \{\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4\}\,,\\ \Lambda_{abcd} &= \Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}} \ni \{\Lambda_5, \Lambda_6, \Lambda_7\}\,,\\ \Lambda_{abc\bar{d}} &= \Lambda_{a\bar{b}\bar{c}\bar{d}} \ni \{\Lambda_8, \Lambda_9, \Lambda_{10}, \Lambda_{11}\}\,,\end{split}$$

where

$$\begin{split} M^2 &\equiv M_{1\bar{1}}^2 = M_{2\bar{2}}^2, & \bar{M}^2 \equiv M_{11}^2 = -M_{22}^2, \\ \Lambda_1 &\equiv \Lambda_{1111} = \Lambda_{2222}, & \Lambda_2 \equiv \Lambda_{12\bar{1}\bar{2}}, & \Lambda_3 \equiv \Lambda_{11\bar{2}\bar{2}}, \\ \Lambda_4 &\equiv \Lambda_{11\bar{1}\bar{2}} = -\Lambda_{12\bar{2}\bar{2}}, & \Lambda_5 \equiv \Lambda_{1122}, & \Lambda_6 \equiv \Lambda_{1111} = \Lambda_{2222}, \\ \Lambda_7 &\equiv \Lambda_{1112} = -\Lambda_{1222}, & \Lambda_8 \equiv \Lambda_{112\bar{1}} = \Lambda_{122\bar{2}}, & \Lambda_9 \equiv \Lambda_{111\bar{1}} = -\Lambda_{222\bar{2}}, \\ \Lambda_{10} &\equiv \Lambda_{112\bar{2}} = -\Lambda_{122\bar{1}}, & \Lambda_{11} \equiv \Lambda_{111\bar{2}} = \Lambda_{222\bar{1}}. \end{split}$$

#### Realification and Complexification summarized

These terms are inspired by their usage in Lie algebra theory.

as applied to scalar field theory	Lie algebra example		
theory of one complex scalar $\Phi$	$\mathfrak{sl}(2,\mathbb{C})$		
↓ realify	↓ realify		
theory of two real scalars $arphi_1$ , $arphi_2$	$\mathfrak{sl}(2,\mathbb{C})_{\mathbb{R}}\cong\mathfrak{so}(3,1)$		
	$\downarrow$ complexify		
theory of two complex scalars $\Phi_1$ , $\Phi_2$	$\mathfrak{sl}(2,\mathbb{C})\oplus\mathfrak{sl}(2,\mathbb{C})\cong\mathfrak{so}(4,\mathbb{C})$		

<u>Note:</u> The Lie algebra  $\mathfrak{sl}(2,\mathbb{C})$  is defined to be the set of complex traceless  $2 \times 2$  matrices. Any element of  $\mathfrak{sl}(2,\mathbb{C})$  is given by *complex* linear combination of the three generators,  $\left\{\begin{pmatrix}0&1\\0&0\end{pmatrix},\begin{pmatrix}0&0\\1&0\end{pmatrix},\begin{pmatrix}1&0\\0&-1\end{pmatrix}\right\}$ . The realified version, denoted by  $\mathfrak{sl}(2,\mathbb{C})_{\mathbb{R}}$ , consists of *real* linear combinations of the six generators,  $\left\{\begin{pmatrix}0&1\\0&0\end{pmatrix},\begin{pmatrix}0&0\\1&0\end{pmatrix},\begin{pmatrix}1&0\\0&-1\end{pmatrix},\begin{pmatrix}0&i\\0&-1\end{pmatrix},\begin{pmatrix}0&0\\0&0\end{pmatrix},\begin{pmatrix}i&0\\0&-i\end{pmatrix}\right\}$ . The realified version written in a different way.

#### $\beta$ functions of the complexified theory

To use the formulae previously given for a real scalar field theory, we could carry out the realification procedure one more time by defining  $\Phi_1 = (\varphi_1 + i\varphi_2)/\sqrt{2}$  and  $\Phi_2 = (\varphi_3 + i\varphi_4)/\sqrt{2}$ . From the resulting  $\beta$  functions, we have derived the corresponding formulae expressed in terms of the complex parameters of  $\mathcal{V}(\Phi_1, \Phi_2)$ . At one-loop, we find:

$$\begin{split} \beta_{M^2_{\bar{a}\bar{b}}} &= 4M^2_{\bar{c}\bar{d}}\Lambda_{cd\bar{a}\bar{b}} + 24M^2_{cd}\Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}} + 6M^2_{e\bar{d}}\Lambda_{d\bar{a}\bar{b}\bar{e}} \,, \\ \beta_{M^2_{a\bar{b}}} &= 12M^2_{cd}\Lambda_{a\bar{b}\bar{c}\bar{d}} + 12M^2_{\bar{c}\bar{d}}\Lambda_{acd\bar{b}} + 8M^2_{d\bar{e}}\Lambda_{ae\bar{b}\bar{d}} \,. \end{split}$$

Apart from the numerical coefficients, the form of these equations is fixed by the index structure of the various terms.

Given that a symmetry of the complexified model imposes the condition  $M_{11}^2 = -M_{22}^2$ , we can write the parameter relation abstractly as

 $c_{ab}M_{\bar{a}\bar{b}}^2 = 0$ , where  $c_{11} = c_{22} = 1$  and  $c_{12} = c_{21} = 0$ .

The symmetry guarantees that

$$c_{ab}\beta_{M^2_{\bar{a}\bar{b}}}\big|_{\rm sym} = c_{ab}\big[4M^2_{\bar{c}\bar{d}}\Lambda_{cd\bar{a}\bar{b}} + 24M^2_{cd}\Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}} + 6M^2_{e\bar{d}}\Lambda_{d\bar{a}\bar{b}\bar{e}}\big]\big|_{\rm sym} = 0\,,$$

where "sym" indicates that the parameter relations of the complexified theory have been applied. But the three quantities in the middle expression above are linearly independent tensors. Thus, each of these quantities must separately vanish.

We conclude that

$$\begin{aligned} c_{ab} M_{\bar{c}\bar{d}}^2 \Lambda_{cd\bar{a}\bar{b}} \Big|_{\rm sym} &= 0 \,, \\ c_{ab} M_{cd}^2 \Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}} \Big|_{\rm sym} &= 0 \,, \\ c_{ab} M_{e\bar{d}}^2 \Lambda_{d\bar{a}\bar{b}\bar{e}} \Big|_{\rm sym} &= 0 \,. \end{aligned}$$

Compare the result in red font to the vanishing of the one-loop beta function of the original toy model of two real scalar fields:

$$\beta_{c_{ij}m_{ij}^2}\big|_{\mathrm{sym}} = c_{ij}m_{k\ell}^2\lambda_{ijk\ell}\big|_{\mathrm{sym}} = 0\,,$$

with  $c_{11} = c_{22} = 1$  and  $c_{12} = c_{21} = 0$  and "sym" indicates that the parameter relations of the real theory have been applied.

Since  $\Lambda_{abcd}$  and  $\Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}}$  are numerically equal (due to CP symmetry), the two equations in red are algebraically identical.

What happens at two-loop order? Recall that

$$\beta_{m_{ij}^2}^{II} = \frac{1}{12} \left( \lambda_{ik\ell m} \lambda_{nk\ell m} m_{nj}^2 + \lambda_{jk\ell m} \lambda_{nk\ell m} m_{ni}^2 \right) - 2m_{k\ell}^2 \lambda_{ikmn} \lambda_{j\ell mn}$$

In the original real scalar theory, if the parameter relation  $c_{ij}m_{ij}^2$  is RG-stable, then

$$c_{ij} \left( \lambda_{ik\ell m} \lambda_{nk\ell m} m_{nj}^2 + \lambda_{jk\ell m} \lambda_{nk\ell m} m_{ni}^2 \right) \Big|_{\text{sym}} = 0,$$
$$c_{ij} m_{k\ell}^2 \lambda_{ikmn} \lambda_{j\ell mn} \Big|_{\text{sym}} = 0.$$

The corresponding equations for  $\beta_{M_{a\bar{b}}^2}^{II}$  and  $\beta_{M_{a\bar{b}}^2}^{II}$  are more complicated, but the index structure fixes the possible terms that can appear. As before, the terms with different index structures must separately vanish.

Due to the symmetry imposed parameter relation  $c_{ab}M^2_{\bar{a}\bar{b}} = 0$ , we find that among the relations obtained from  $\beta^{II}_{M^2_{\bar{a}\bar{b}}}$ ,

$$c_{ab} \left( \Lambda_{\bar{a}\bar{d}\bar{e}\bar{f}} \Lambda_{cdef} M_{\bar{c}\bar{b}}^2 + \Lambda_{\bar{b}\bar{d}\bar{e}\bar{f}} \Lambda_{cdef} M_{\bar{c}\bar{a}}^2 \right) \Big|_{\text{sym}} = 0 ,$$
$$c_{ab} M_{cd}^2 \Lambda_{\bar{a}\bar{c}\bar{e}\bar{f}} \Lambda_{ef\bar{d}\bar{b}} \Big|_{\text{sym}} = 0 .$$

CP symmetry yields  $\Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}} = \Lambda_{abcd}$  and  $M_{\bar{a}\bar{b}}^2 = M_{ab}^2$ . The two displayed equations above hold for any choice of  $\Lambda_{abcd}$  and  $\Lambda_{ab\bar{c}\bar{d}}$ , subject to the parameter relations of the complexified theory:

$$\begin{split} \Lambda_{1111} &= \Lambda_{2222} , \qquad \Lambda_{1112} = -\Lambda_{1222} , \\ \Lambda_{11\bar{1}\bar{1}} &= \Lambda_{22\bar{2}\bar{2}} , \qquad \Lambda_{11\bar{1}\bar{2}} = -\Lambda_{12\bar{2}\bar{2}}^* , \qquad \Lambda_{11\bar{2}\bar{2}} = \Lambda_{11\bar{2}\bar{2}}^* , \end{split}$$

$$c_{ab} \left( \Lambda_{\bar{a}\bar{d}\bar{e}\bar{f}} \Lambda_{cdef} M_{\bar{c}\bar{b}}^2 + \Lambda_{\bar{b}\bar{d}\bar{e}\bar{f}} \Lambda_{cdef} M_{\bar{c}\bar{a}}^2 \right) \Big|_{\text{sym}} = 0 ,$$
$$c_{ab} M_{cd}^2 \Lambda_{\bar{a}\bar{c}\bar{e}\bar{f}} \Lambda_{ef\bar{d}\bar{b}} \Big|_{\text{sym}} = 0 .$$

Since  $\Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}}$  and  $\Lambda_{ab\bar{c}\bar{d}}$  are independent tensors (with compatible relations imposed by the symmetries), the above equations must hold if we numerically set  $\Lambda_{ab\bar{c}\bar{d}} = \Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}}$ . With this choice, the equations above are algebraically equivalent to the corresponding equations of the original real scalar field theory!

Moreover, the same argument extends to arbitrary order in perturbation theory.

## The origin of RG-stable parameter relations

Schematically, 
$$\beta_{c_{ab}M^2_{\bar{a}\bar{b}}} = c_{ab}\sum_k f_k(M^2,\Lambda)_{\bar{a}\bar{b}}.$$

Each term in the sum contains one factor of  $M^2$  and n factors of  $\Lambda$  at order n, where  $M^2$  can have index structure cd,  $cd\bar{d}$ , or cd, and  $\Lambda$  can have index structure cdef,  $cde\bar{f}$ ,  $cd\bar{e}\bar{f}$ ,  $cd\bar{e}\bar{f}$ , or  $cd\bar{e}\bar{f}$ . The indices must combine (including Kronecker deltas) such that the index structure of the  $f_k$  is  $\bar{a}\bar{b}$ . Then,

$$\beta_{c_{ab}M_{\bar{a}\bar{b}}^2}\big|_{\rm sym} = 0 \qquad \Longrightarrow \qquad c_{ab}f_k(M^2,\Lambda)_{\bar{a}\bar{b}}\big|_{\rm sym} = 0\,,$$

for each k separately.

There will always be at least one value of k where  $f_k(M^2, \Lambda)$ depends on tensors with an even number of unbarred and barred indices, respectively. Since  $\Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}}$  and  $\Lambda_{ab\bar{c}\bar{d}}$  are independent, we can numerically set  $\Lambda_{ab\bar{c}\bar{d}} = \Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}}$ . Thus, for some value of k,

$$c_{ab}f_k(M^2,\Lambda)_{\bar{a}\bar{b}}\big|_{\mathrm{sym}} = 0,$$

where the distinction between unbarred and barred indices can be neglected.<sup>13</sup> This equation will be algebraically equivalent to the corresponding equation of the original real scalar field theory.

That is, the RG-stability of the parameter relations of the original real scalar field theory is inherited from the symmetry of the corresponding complexified theory.

<sup>&</sup>lt;sup>13</sup>Having imposed CP symmetry, all scalar potential parameters are real.

# Future directions

1. A recipe to create real scalar field theories with RG-stable parameter relations in absence of a symmetry:

- Start with a theory of n complex scalars Φ<sub>a</sub> with RG-stable parameter relations due to some HF symmetry [which is a subgroup of U(n), the symmetry group of L<sub>KE</sub>].
- Impose CP to ensure the reality of all scalar potential parameters.
- Retain only the terms of the scalar potential that are holomorphic in the complex fields, i.e., of the form  $M_{\bar{a}\bar{b}}^2 \Phi_a \Phi_b + \Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}} \Phi_a \Phi_b \Phi_c \Phi_d.$

- Construct the corresponding theory of *n* real scalars with the following recipe.
  - Replace the  $\Phi_a$  with n real scalar fields  $\varphi_a$ .
  - Replace  $\mathscr{L}_{KE}$  with a canonically normalized kinetic energy term for the real scalar theory [with symmetry group O(n)].
- If the HF symmetry of the complex scalar field theory cannot be embedded in O(n), then this HF symmetry will not survive as a legitimate symmetry of the real scalar field theory.

However, the symmetry-imposed parameter relations satisfied by  $M_{\bar{a}\bar{b}}$  and  $\Lambda_{\bar{a}\bar{b}\bar{c}\bar{d}}$  are now parameter relations of the same form in the resulting real scalar field theory. These parameter relations are RG-stable due to the symmetries of the complexified theory.

- 2. Applying the results of this talk to the GOOFy symmetries of the 2HDM.
- Starting from the 2HDM Lagrangian written in terms of complex doublets, perform the realification procedure to rewrite the theory in terms of eight real scalar fields.

$$\Phi_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} \varphi_1 + i\varphi_2 \\ \varphi_3 + i\varphi_4 \end{pmatrix}, \qquad \Phi_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} \varphi_5 + i\varphi_6 \\ \varphi_7 + i\varphi_8 \end{pmatrix},$$

• The GOOFy symmetries take the form

$$\begin{split} \varphi_1 &\to i\varphi_6 \,, \qquad \varphi_2 \to i\varphi_5 \,, \qquad \varphi_3 \to i\varphi_8 \,, \qquad \varphi_4 \to i\varphi_7 \,, \\ \varphi_5 &\to -i\varphi_2 \,, \quad \varphi_6 \to -i\varphi_1 \,, \quad \varphi_7 \to -i\varphi_4 \,, \quad \varphi_8 \to -i\varphi_3 \,. \end{split}$$

- Complexify the theory by promoting the  $\varphi_i$  to eight complex scalar fields  $\Phi_a$ .
- Impose CP symmetry to ensure that all scalar potential parameters are real.
- Verify that the RG-stability of the 2HDM with  $m_{11}^2 = -m_{22}^2$ ;  $\lambda_1 = \lambda_2$ ;  $\lambda_6 = -\lambda_7$  can be explained by the symmetries of the complexified theory. This step will require an extension of the techniques of this work to the gauge and Yukawa sectors.

3. Are there more sophisticated ideas that can be applied to provide a deeper understanding of the phenomena explored in this work? (Outer automorphisms? Generalized symmetries?)<sup>14</sup>

<sup>&</sup>lt;sup>14</sup>See, e.g., Andreas Trautner, *Goofy is the new Normal*, arXiv:2505.00099 [hep-ph].

# Backup slides

References to beta functions of real scalar field theories

T.P. Cheng, E. Eichten and L.F. Li, Phys. Rev. D 9 (1974) 2259.

M.E. Machacek and M.T. Vaughn, Nucl. Phys. B 249 (1985) 70.

M.x. Luo, H.w. Wang and Y. Xiao, Phys. Rev. D **67** (2003) 065019, arXiv:hep-ph/0211440 [hep-ph].

I. Schienbein, F. Staub, T. Steudtner and K. Svirina, Nucl. Phys. B **939** (2019) 1 [erratum: **966** (2021) 115339], arXiv:1809.06797 [hep-ph].

L. Sartore, Phys. Rev. D 102 (2020) 076002, arXiv:2006.12307 [hep-ph].

T. Steudtner, JHEP 12 (2020) 012 (2020), arXiv:2007.06591 [hep-th].

References to beta functions of the 2HDM (up to three loops)

F. Herren, L. Mihaila and M. Steinhauser, Phys. Rev. D **97** (2018) 015016, arXiv:1712.06614 [hep-ph].

A.V. Bednyakov, JHEP 11 (2018) 154, arXiv:1809.04527 [hep-ph].