

70 years of Howie

The last \sim 23 Years:

Two papers and many² workshops

What you really want to see: photos

→ I found \sim 150 photos

→ will show only a fraction ;-)

...surprisingly few photos in which Howie is working ...

Let's start with the papers:

Inspire search: a haber and heinemeyer → 23 results

The first and the last:

Reconciling the Two-Loop Diagrammatic and Effective Field Theory Computations of the Mass of the Lightest \mathcal{CP} -even Higgs Boson in the MSSM

M. Carena^{§,†}, H.E. Haber[#], S. Heinemeyer[‡],

W. Hollik[¶], C.E.M. Wagner^{†,*,‡} and G. Weiglein[†]

[§] FERMILAB, Batavia, IL 60510-0500 USA

[†] CERN, TH Division, CH-1211 Geneva 23, Switzerland

[#] Santa Cruz Inst. for Part. Phys., Univ. of California, Santa Cruz, CA 95064 USA

[‡] DESY Theorie, Notkestrasse 85, 22603 Hamburg, Germany

[¶] Institut für Theoretische Physik, Univ. of Karlsruhe, 76128 Karlsruhe, Germany

^{*} High Energy Physics Division, Argonne National Lab., Argonne, IL 60439 USA

[‡] Enrico Fermi Institute, Univ. of Chicago, 5640 Ellis, Chicago, IL 60637 USA

Abstract

The mass of the lightest \mathcal{CP} -even Higgs boson of the minimal supersymmetric extension of the Standard Model (MSSM) has previously been computed including $\mathcal{O}(\alpha\alpha_s)$ two-loop contributions by an on-shell diagrammatic method, while approximate analytic results have also been obtained via renormalization-group-improved effective potential and effective field theory techniques. Initial comparisons of the corresponding two-loop results revealed an apparent discrepancy between terms that depend logarithmically on the supersymmetry-breaking scale, and different dependences of the non-logarithmic terms on the squark mixing parameter, X_t . In this paper, we determine the origin of these differences as a consequence of different renormalization schemes in which both

Higgs-mass predictions in the MSSM and beyond

P. Slavich^a and S. Heinemeyer^{b,c,d} (eds.),

E. Bagnaschi^e, H. Bahl^f, M. Goodsell^a, H.E. Haber^g, T. Hahn^h, R. Harlanderⁱ,
W. Hollik^h, G. Lee^{j,k,l}, M. Mühlleitner^m, S. Paßehrⁱ, H. Rzehakⁿ, D. Stöckinger^o,

A. Voigt^p, C.E.M. Wagner^{q,r,s} and G. Weiglein^f,

B.C. Allanach^t, T. Biekötter^f, S. Borowka^{u,‡}, J. Braathen^f, M. Carena^{r,s,v},
T.N. Dao^w, G. Degrandi^x, F. Domingo^y, P. Drechsel^{f,‡}, U. Ellwanger^z, M. Gabelmann^m,
R. Gröber^{aa}, J. Klappertⁱ, T. Kwasnitza^o, D. Meuser^f, L. Mihaila^{bb,‡}, N. Murphy^{cc,‡},
K. Nickel^{y,‡}, W. Porod^{dd}, E.A. Reyes Rojas^{ee}, I. Sobolev^f and F. Staub^{m,‡}

Predictions for the Higgs masses are a distinctive feature of supersymmetric extensions of the Standard Model, where they play a crucial role in constraining the parameter space. The discovery of a Higgs boson and the remarkably precise measurement of its mass at the LHC have spurred new efforts aimed at improving the accuracy of the theoretical predictions for the Higgs masses in supersymmetric models. The “*Precision SUSY Higgs Mass Calculation Initiative*” (KUTS) was launched in 2014 to provide a forum for discussions between the different groups involved in these efforts. This report aims to present a comprehensive overview of the current status of Higgs-mass calculations in supersymmetric models, to document the many advances that were achieved in recent years and were discussed during the KUTS meetings, and to outline the prospects for future improvements in these calculations.

They have more in common than one may think ...

Reconciling the Two-Loop Diagrammatic and Effective Field Theory Computations of the Mass of the Lightest \mathcal{CP} -even Higgs Boson in the MSSM

M. Carena ^{§,†}, H.E. Haber [#], S. Heinemeyer [‡],

W. Hollik [¶], C.E.M. Wagner ^{†,*,‡} and G. Weiglein [†]

[§] FERMILAB, Batavia, IL 60510-0500 USA

[†] CERN, TH Division, CH-1211 Geneva 23, Switzerland

[#] Santa Cruz Inst. for Part. Phys., Univ. of California, Santa Cruz, CA 95064 USA

[‡] DESY Theorie, Notkestrasse 85, 22603 Hamburg, Germany

[¶] Institut für Theoretische Physik, Univ. of Karlsruhe, 76128 Karlsruhe, Germany

^{*} High Energy Physics Division, Argonne National Lab., Argonne, IL 60439 USA

[‡] Enrico Fermi Institute, Univ. of Chicago, 5640 Ellis, Chicago, IL 60637 USA

Abstract

The mass of the lightest \mathcal{CP} -even Higgs boson of the minimal supersymmetric extension of the Standard Model (MSSM) has previously been computed including $\mathcal{O}(\alpha\alpha_s)$ two-loop contributions by an on-shell diagrammatic method, while approximate analytic results have also been obtained via renormalization-group-improved effective potential and effective field theory techniques. Initial comparisons of the corresponding two-loop results revealed an apparent discrepancy between terms that depend logarithmically on the supersymmetry-breaking scale, and different dependences of the non-logarithmic terms on the squark mixing parameter, X_t . In this paper, we determine the origin of these differences as a consequence of different renormalization schemes in which both

⇒ note the arXiv number!

⇒ the second paper in the new millenium

(modulo 2000 vs. 2001)

Now for the workshops :-)

The oldest I could find: **LCWS '05** (San Francisco)



Howie is a frequent participant in the “Higgs Days at Santander”

Q: What is Howies HDays number?

Howie is a frequent participant in the “Higgs Days at Santander”

Q: What is Howies HDays number?

A: 7 (among the top ~ 10 :-)

⇒ a few impression ...

... with different people and different moods

HDays10: only the group photo



HDays12: arguing ...



HDays12 at the workshop dinner:



Not sure what he is telling about Carlos ...

HDays13: at the lighthouse



HDays13: working at the workshop dinner



HDays14: in the 1-star restaurant



HDays16: reception at the roof top



HDays16: the summer with the tooth problem ;-)



HDays17: tooth is in best shape again :-)



HDays19: again at the workshop dinner



⇒ Is he ever working?

HDays19: finally working!



... on cartoons?!

HDays20: even during the pandemic!



But there were many other workshops!

- Higgs Hunting (Paris)
- HPNP (Toyama, Osaka)
- SUSY (everywhere)
- Multi-Higgs (Lisbon)
- Multi-Higgs (Taiwan)
- LC workshops (everywhere)
- Snowmass (Seattle ;-)
- . . .

⇒ a few more impressions

HiggsHunting '16: Reisaburo organizes a dinner



HiggsHunting '18: high above Paris




HPNP '15: It was really, really cold



HPNP '15: the best after-dinner speech ever



HPNP '17: finally working again



HPNP '17: and back to the workshop dinner ;-)




HPNP '19: with the organizer!



HPNP '19: and with a nice participant!



Snowmass in Seattle 2013:



Out of the many SUSY conferences just one: 2015




Howie showed me how to drive and eat at the same time

Out of the many SUSY conferences just one: 2015



But it was worth the risk!

Multi-Higgs in Lisbon '18: Howie as M.C. Escher



Multi-Higgs in Lisbon '18: very large Howie



Multi-Higgs in Lisbon '18: very small Howie



And finally: one photo at home :-)

And finally: one photo at home :-)




And very finally: a photo from SUSY'09 ...



... that introduces the second paper

Howie as a model for the poster of the KUTS workshop series:



Katharsis of Ultimate Theory Standards

11th meeting: 20-22 November 2019 (MPI Munich)

Precise Calculation of

(N) ~~100~~

Higgs Boson masses

Local organizers: T. Hahn, W. Hollik

Organized by:
M. Carena, H. Haber
R. Harlander, S. Heinemeyer
W. Hollik, P. Slavich, G. Weiglein

Higgs-mass predictions in the MSSM and beyond

P. Slavich^a and S. Heinemeyer^{b,c,d} (eds.),

E. Bagnaschi^e, H. Bahl^f, M. Goodsell^a, H.E. Haber^g, T. Hahn^h, R. Harlanderⁱ,
W. Hollik^h, G. Lee^{j,k,l}, M. Mühlleitner^m, S. Paßehrⁱ, H. Rzehakⁿ, D. Stöckinger^o,
A. Voigt^p, C.E.M. Wagner^{q,r,s} and G. Weiglein^f,

B.C. Allanach^t, T. Biekötter^f, S. Borowka^{u‡}, J. Braathen^f, M. Carena^{r,s,v},
T.N. Dao^w, G. Degrandi^x, F. Domingo^y, P. Drechsel^{f‡}, U. Ellwanger^z, M. Gabelmann^m,
R. Gröber^{aa}, J. Klappertⁱ, T. Kwasnitza^o, D. Meuser^f, L. Mihaila^{bb‡}, N. Murphy^{cc‡},
K. Nickel^{y‡}, W. Porod^{dd}, E.A. Reyes Rojas^{ee}, I. Sobolev^f and F. Staub^{m‡}

Predictions for the Higgs masses are a distinctive feature of supersymmetric extensions of the Standard Model, where they play a crucial role in constraining the parameter space. The discovery of a Higgs boson and the remarkably precise measurement of its mass at the LHC have spurred new efforts aimed at improving the accuracy of the theoretical predictions for the Higgs masses in supersymmetric models. The “*Precision SUSY Higgs Mass Calculation Initiative*” (KUTS) was launched in 2014 to provide a forum for discussions between the different groups involved in these efforts. This report aims to present a comprehensive overview of the current status of Higgs-mass calculations in supersymmetric models, to document the many advances that were achieved in recent years and were discussed during the KUTS meetings, and to outline the prospects for future improvements in these calculations.

arXiv:2012.15629v3 [hep-ph] 17 May 2021

In 2020 we got it out juuuuust in time :-)

Happy Birthday, Howie!!



Looking forward to the next meeting in 3D!